

# Quantification of silkworm coactivator of MBF1 mRNA by SYBR Green I real-time RT-PCR reveals tissue- and stage-specific transcription levels

Guang-li Li · Bhaskar Roy · Xing-hua Li ·  
Wan-fu Yue · Xiao-feng Wu · Jian-mei Liu ·  
Chuan-xi Zhang · Yun-gen Miao

Received: 21 February 2008 / Accepted: 18 June 2008 / Published online: 9 July 2008  
© Springer Science+Business Media B.V. 2008

**Abstract** Transcriptional coactivators play a crucial role in gene transcription and expression. Multiprotein bridging factor 1 (MBF1) is a transcriptional coactivator necessary for transcriptional activation caused by DNA-binding activators, such as FTZ-F1 and GCN4. Until now, very few studies have been reported in the silkworm. We selected the *Bombyx mori* because it is a model insect and acts as an economic animal for silk industry. In this study, we conducted the quantitative analysis of MBF1 mRNA in silkworm *B. mori* L. with actin (A3) as internal standard by means of SYBR Green I real-time RT-PCR method. The total RNA was extracted from the silk gland, epidermis, fat body, and midguts of the fifth instar *B. mori* larvae. The mRNA was reverse transcribed, and the cDNA fragments of MBF1 mRNA and actin gene were amplified by RT-PCR using specific primers. MBF1 mRNA expression in different tissues of silkworm *B. mori* L. was quantified using standardized SYBR Green I RT-PCR. The results suggested MBF1 gene was expressed in all investigated organs but highly expressed in the silk gland, showing its relation to biosynthesis of silk proteins.

**Keywords** RT-PCR · SYBR Green I real-time RT-PCR · Multiprotein Bridging Factor 1 (MBF1) gene · Silkworm *Bombyx mori* L.

## Introduction

Many *cis*-regulatory elements on DNA and regulatory proteins that bind to them in a sequence-specific manner are involved in the control of gene expression [1, 2]. Subsequent studies have shown that non-DNA-binding factors termed coactivators and mediators play an important role by interconnecting the DNA-binding regulators and basal transcription machinery [3–5]. Proper combination of the *cis* elements, DNA-binding regulators, and coactivators appears to be critical for temporal and spatial control of gene expression [6].

MBF1 has been identified as a transcriptionally coactive factor from yeast, plant, and animal cells [7, 8]. MBF1 bridges between the TATA box-binding protein (TBP) and the *Drosophila melanogaster* nuclear hormone receptor FTZ-F1 (Fushi tarazu transcriptional factor 1) or its silkworm counterpart BmFTZ-F1 [9, 10]. Tethering of the positive cofactor MBF2 to a FTZ-F1-binding site through FTZ-F1 and MBF1 is essential for the binding site-dependent activation of transcription. A homology search in the databases revealed that the deduced amino acid sequence of MBF1 is conserved across species from yeast to human.

The silk production by *Bombyx mori* makes it one of the most economic important insects in the world. It is employed for silk production. It is also one of the best-characterized models for biochemical, molecular, genetic, and genomic studies of the order Lepidoptera because of its large size, complex metabolism, and the abundance of mutants [11, 12]. The ability to study molting and morphogenesis demonstrates how *B. mori* is advantageous because precisely staged samples are available according to morphological changes of the spiracle [13]. Furthermore, the fifth instar is a transition period for the metamorphosis from larvae to pupa, and for biosynthesizing and spinning

G.-l. Li · B. Roy · X.-h. Li · W.-f. Yue ·  
X.-f. Wu · J.-m. Liu · Y.-g. Miao (✉)  
College of Animal Sciences, Institute of Sericulture  
and Apiculture, Zhejiang University, Hangzhou 310029,  
People's Republic of China  
e-mail: miaoyg@zju.edu.cn

C.-x. Zhang  
Institute of Insect Sciences, Zhejiang University,  
Hangzhou 310029, People's Republic of China

silk proteins. The molecular mechanisms involved in this short period are still not well-understood.

Multiprotein bridging factor 1 (MBF1) is a transcriptional coactivator. But very few researches in silkworm were reported. It is meaningful to realize its function of MBF1 gene because silkworm is a model insect and its unique silk synthesis character. This study describes expression patterns of MBF1 mRNA in the silk gland, epidermis, fat body, and midgut of the fifth instar larvae during the development of *B. mori*. We conducted the quantitative analysis of MBF1 mRNA with actin (A3) as the internal standard through SYBR Green I real-time RT-PCR method. The results showed MBF1 gene was expressed in all investigated organs but highly expressed in the silk gland. This suggests that MBF1 plays an important role in the development of the silkworm as a tissue-specific and stage-specific coactivator, showing its relation to biosynthesis of silk proteins.

## Materials and methods

### Experimental animals

The silkworm larvae of strain P50 were reared on mulberry leaves at 25°C under a 16 h light, 8 h dark photoperiod. The silk gland, epidermis, fat body and midgut of the fifth instar larvae (10 larvae as a sample) were dissected for analysis of developmental expression patterns.

### Chemistry reagents

TRIZOL, pUCm-T vector, EZ spin column plasmid maxi-preps kit (Invitrogen, USA), fast ligation kit and hot start fluorescent PCR core reagent kits (SYBR Green I) were bought from Bio Basic Inc., Canada. MMLV first-strand cDNA synthesis kit was purchased from Sangon Company, Shanghai, China.

### RNA extraction and cDNA synthesis

The total RNA was extracted from samples using TRIZOL reagent. The RNA was used as a template for first-strand cDNA synthesis using M-MLV reverse transcriptase with oligo dT<sub>18</sub> primer under reaction conditions of 5 min at 70°C, 10 min at 37°C, 60 min at 37°C, 10 min at 70°C. For gene fragment amplification, two primers were designed as MBF1's forward primer 5'-AAggtccATGTCTGACTGGG ATACAGT, and reverse primer 5'-ACAAagcttTTATTTCT GTCCGCCAGGAG; Actin's forward primer 5'-GCGCG GCTACTCGTTCCTACTACC, and reverse primer 5'-GGA TGTCCACGTCGCACTTCA.

The PCR was conducted in 20 µl system containing 13.1 µl double distilled water, 2 µl 10 × PCR buffer,

0.2 µl 25 mmol/l dNTP mix, 1 µl 10 mmol/l primer I, 1 µl 10 mmol/l primer II, 1.2 µl 25 mmol/l MgCl<sub>2</sub>, 0.5 µl Taq DNA polymerase, 1 µl cDNA. The condition of reaction was set as 50 s at 94°C, 50 s at 60°C, 1 min at 72°C and final extension for 10 min at 72°C.

### Cloning the PCR product and sequencing

The RT-PCR generated products were T/A cloned into pUCm 18-T vector and then sequenced.

### Comparison of MBF1 transcription by semi-quantitative RT-PCR

The comparison of MBF1 transcription by semi-quantitative RT-PCR was performed according to a modified procedure of previously described protocol [14]. Actin gene, a house-keeping gene, was used as a reference gene to allow for normalization by visual inspection of mRNA levels. The quantity of products was screened in agarose gels. All experiments were repeated three times.

### SYBR real-time quantitative PCR

PCR amplifications were performed using a Bio-RAD IQ<sup>TM</sup>5 Multicolor real-time PCR detection system. The PCR was conducted in 25 µl system containing 9.5 µl ddH<sub>2</sub>O, 12.5 µl Hotstart Fluo-PCR mix, 1 µl 10 mmol/l primer I, 1 µl 10 mmol/l Primer II, 1 µl cDNA. The PCR protocol was done by 30 s denaturation at 94°C, 30 s annealing at 60°C, and 30 s elongation at 72°C in 35 cycles. Fluorescence was detected at the end of every 72°C extension phase. To exclude the contamination of unspecific PCR products such as primer dimers, melting curve analysis was applied to all final PCR products after the cycling protocol.

### Construction of the standard curve of the MBF1 and actin gene

To determine the copy number of the target transcript, the cloned plasmid DNA for MBF1 and actin were used to generate the standard curve. The plasmid DNA was purified using alkaline prep. Plasmids contained the cDNA inserts encoding the respective PCR products in a pUCm-T vector. The cloned plasmid DNA were consecutively diluted every tenfold at a range of 10<sup>7</sup>–10<sup>3</sup> copies.

### Comparison of MBF1 transcription by SYBR Green I RT-PCR

This was performed similar to the standard curve. Each cDNA from the silk gland, epidermis, fat body, and midgut of the first, fourth and eighth day of the fifth instar larvae

were run in triplicates. The threshold cycle ( $C_t$ ) values were averaged from each reaction. Actin gene isolated in this study was used as a reference gene to normalize mRNA levels.

#### Data analysis

The data were expressed as mean  $\pm$  SD, and compared statistically by *t*-test,  $P < 0.05$  being considered significant.

## Results

### Clone and sequence of the MBF1 gene

The total RNA was extracted from samples using the TRIZOL reagent. The RNA was used as the template for first-strand cDNA synthesis using M-MLV reverse transcriptase with oligo dT<sub>18</sub> primer. The PCR was conducted using special primers for MBF1 and actin gene amplification. The electrophoresis bands of PCR product is shown in Fig. 1.

The RT-PCR generated products were T/A cloned into pUCm 18-T vector and then sequenced. The sequence of MBF1 cDNA is shown in Fig. 2. GenBank Blast search revealed that the resulting nucleotide sequences of MBF1 and actin gene shared the same identity with silkworm *B. mori*. It contained a single open reading frame of 146 amino acids. The deduced amino acid sequence of the factor predicts a basic protein (*pI* 10.7) with a molecular mass of 16.2 kDa.

### Expression of MBF1 mRNA in organs with semi-quantitative RT-PCR

We used RT-PCR method to amplify MBF1 and actin gene from the silk gland, epidermis, fat body, and midgut. The PCR was carried out for 28 cycles for MBF1 gene and

30 cycles for actin gene. The electrophoresis bands of PCR product is shown in Fig. 3.


As Fig. 3 shows, the highest expression level of MBF1 gene was detected in the silk gland. The accumulation of MBF1 mRNA in the silk gland reached its peak at the first day of the fifth instar larvae and then decreased significantly. The expressions of the MBF1 gene in the epidermis, fat body, and midgut were lower than that of the silk gland. In the midgut, the expression level of MBF1 gene has demonstrated an increasing trend with the development of larvae.

### Standard curve analysis for SYBR Green I real-time quantitative PCR

Standard curves for each MBF1 and actin gene were obtained using standard plasmids that encodes the PCR products. The quantitative nature is demonstrated by the linear relationship between the log of the template copy number and  $C_t$  value. We also confirmed the linear quantitative relationship between the  $C_t$  of each gene and the log of the t concentrations of consecutively diluted DRG samples. A range with a correlation value (*R*) greater than 0.99 was described as quantitative (Fig. 4a, b). Melting curve analysis revealed that PCR amplified a single desired product (Fig. 4c). When comparing the quantity of target gene samples, actin was used as a housekeeping gene to correct the total non-degraded RNA amount among MBF1 samples. We examined the expression of actin using quantitative real-time PCR analysis (Fig. 5a–c).

### Expression pattern of MBF1 mRNA in silkworm *B. mori* L. organs

SYBR Green I RT-PCR protocols were established for MBF1 and actin gene. All organs analyzed by SYBR Green I RT-PCR expressed MBF1 mRNA. The ratio between MBF1 and actin transcript numbers ranged from 0.03 in the



**Fig. 1** The electrophoresis bands of PCR product. (a) is MBF1 gene, a 458 bp band was identified in agarose gel electrophoresis. (b) is actin gene, a 284 bp band was identified


```

1 atgtctgactgggatacagttacaattcttcgcaagaaccacacaaagcttctgctttaaact
  M S D W D T V T I L R K K P P K A S A L K T
67 gaacaagctgtaaatgcagcggcgacaaggtattccggttgatagcaacaaaatattgggct
  E Q A V N A A R R Q G I P V D T Q Q K Y G A
133 ggaactaataaacaacatgttactactaaaaatacagctaaactagacagagaacaagaatta
  G T N K Q H V T T K N T A K L D R E T E E L
199 cgccatgaaaaataaccattgatctgggaaacttataatgcaaggcagacaagctaagggaatg
  R H E K I P L D L G K L I M Q G R Q A K G M
265 agtcaaaaagacctcgccactaaaaatttgtaaaaaccacagattgtaaatgattatgaagctggt
  S Q K D L A T K I C E K P Q I V N D Y E A G
331 cgtggcattccaataacattgttcttggaaaaaattgaaaggccaattggaataaaactctgtgga
  R G I P N N I V L G K I E R A I G I K L R G
397 aaagaagaggccaaccactgcagcctctggcggacagaaaataa
  K E R G Q P L Q P P G G Q K *

```

**Fig. 2** Sequence of MBF1 cDNA. The predicted amino acid sequence of the factor is presented in single-letter code. The asterisk represents the putative stop codon

**Fig. 3** Semi-quantitative RT-PCR analysis of MBF1 transcripts in organs. Agarose gel analyses of MBF1 amplifications in comparison with those of controls (actin). (1) Molting fourth instar; (2) the first day of the fifth instar larval; (3) the third day of the fifth instar larval; (4) the fifth day of the fifth instar larval; (5) the seventh day of the fifth instar larval



midgut to 8.57 in the silk gland of the first day of the fifth instar (Table 1 and Fig. 6). As Fig. 6 shows, the highest expression level of MBF1 gene was detected in the silk gland. The accumulation of MBF1 mRNA in the silk gland reached its peak at the first day of the fifth instar larvae and then decreased. The statistical significance is evidently different between different growths. Based on these results, hypothesize that the silk protein synthesis is activated in the beginning of the fifth instar. The MBF1 gene regulated the procedure in this stage.

Furthermore in the midgut, the expression of MBF1 mRNA has revealed the reverse phenomenon. The expression increased with the development of the fifth instar, which was consistent with the results of semi-quantitative RT-PCR analysis. In the epidermis and fat body, the expression of MBF1 mRNA significantly decreased in the fourth day and reached its peak at the eighth day of the fifth instar larvae (Fig. 7).


## Discussion

MBF1 contains a nuclear export signal (NES)-like sequence in its C-terminal region, which is conserved among eukaryotes [15]. For example, amino acid residues 119–130 of *B. mori* MBF1, LGKIERAIGIKL, and the corresponding region of human MBF1, LGIERAIGLKL [16], are similar to the leucine-rich NES in HIV Revprotein, LPPLERLTL, and protein kinase inhibitor, LALKLAGLDI [17]. During larvae development, BmFTZ-F1, MBF1, and MBF2 form a nuclear complex at the molting stage D3. Transient expression of a reporter gene in S2 cells demonstrated that MBF1 and MBF2 enhance BmFTZ-F1-dependent transcription. These observations support that MBF1 and MBF2 serve as coactivators that mediate BmFTZ-F1-dependent transcriptional activation [18]. MBF1 can significantly stabilize BmFTZ-F1 binding to DNA [10, 19].

Takemaru et al. provides evidence acquired in vitro and in vivo that yeast MBF1 mediates GCN4-dependent transcriptional activation by bridging the DNA-binding region of GCN4 and TBP [8, 20]. These findings indicate that the coactivator MBF1 functions by recruiting TBP to promoters where DNA-binding regulators are bound [21]. Through immuno-precipitation, MBF1 was found to form a ternary complex including MBF1, TBP, and the bZIP protein tracheae defective (TDF)/Apontic. Many data demonstrate a crucial role of MBF1 in the development of the tracheae and central nervous system [22] via a direct interaction with the basic region of *Drosophila* Jun (D-Jun), MBF1 prevents an oxidative modification (S-cystenyl cystenylation) of the critical cysteine and stimulates AP-1 binding to DNA by preserving the redox-sensitive AP-1 activity. It has been shown that human MBF1 stimulates the transcriptional activity of steroidogenic factor 1, a human homolog of fushi tarazu factor 1, which is implicated in steroidogenesis [23]. Plant MBF1 can be rapidly induced by several stresses whereas animal MBF1 was not induced. MBF1 function in plants is controlled on the level of transcriptional induction, but not by nuclear translocation, unlike the case of MBF1 in animals [24].


The sensitivity of SYBR Green I analysis is more than adequate for the majority of research applications, and its quantitative nature is demonstrated by the linear relationship between the log of template concentration and cycle number at which fluorescence rises above baseline. The increased sensitivity and specificity of SYBR Green I real-time assay obtained by increasing the amplification size coupled with the low cost of SYBR Green dye as opposed to TaqMan probes will help to extend the applicability of SYBR Green I real-time PCR in detecting other viral pathogens as well as in cellular gene expression studies.

In this study, we described the expression pattern of MBF1 mRNA using both the semi-quantitative RT-PCR and the SYBR Green I real-time quantitative RT-PCR method. During the larvae stage, MBF1 mRNA was



**Fig. 4** Example of real-time PCR for quantification of MBF1 mRNA. (a) Example of real-time PCR amplification stand curves of MBF1; (b) example of real-time PCR amplification recursive equation obtained by plotting fluorescence data against their cycle number; (c) sample of real-time PCR amplification melting curves. The  $T_m$  is 84°C, the production is specific

constitutively expressed in all investigated organs. It can be considered MBF1 is a very important gene in the growth of *B. mori*. The highest expression level of MBF1 gene was detected in the silk gland, revealing the strong relation with biosynthesis of silk proteins. The accumulation of MBF1 mRNA in the silk gland reached its peak at the first day of the fifth instar larvae and then decreased significantly. This implies high MBF1 activity is necessary for silk protein gene transcription in the beginning of fifth instar, and the MBF1 expression was reduced with the gradual




**Fig. 5** Example of real-time PCR for quantification of actin mRNA. (a) Example of real-time PCR amplification stand curves of actin; (b) example of real-time PCR amplification recursive equation obtained by plotting fluorescence data against their cycle number; (c) sample of real-time PCR amplification melting curves. The  $T_m$  is 84°C, the production is specific

deterioration of the silk gland and ultimately silk protein synthesis ceased at the eighth day of the fifth instar. On the contrary, the expression of MBF1 mRNA reached its peak in epidermis and fat body at the eighth day of the fifth instar larvae, suggesting MBF1 regulated the genes of metamorphosis such as chitinase and fatty acid synthetase. Thus, on the basis of this present research, we conclude that multiprotein bridging factor 1 (MBF1) is required for transcriptional activation of genes during the silkworm development.


**Table 1** Relative quantification of MBF1 mRNA in organs by comparison with those of controls (actin)

Stage	Silk gland	Epidermis	Fat body	Middle gut
First day of the fifth instar	8.5745 ± 0.0306	0.1481 ± 0.0290	0.3051 ± 0.1005	0.0302 ± 0.0061
Fourth day of the fifth instar	5.4731 ± 0.0493**	0.0338 ± 0.0067**	0.0992 ± 0.1056**	0.1124 ± 0.0113*
Eighth day of the fifth instar	1.4881 ± 0.3842**	0.1787 ± 0.0084*	0.4702 ± 0.0691**	0.2116 ± 0.1013**

Unit: MBF1 mRNA copy/actin mRNA copy. Values are shown as the mean ± SD. \*  $P < 0.05$ , \*\*  $P < 0.01$



**Fig. 6** MBF1 mRNA expression in all organs of fifth instar. The levels of each MBF1 mRNA and actin were measured by SYBR Green I real-time RT-PCR. Silk gland's MBF1 mRNA expression was significantly higher than that of other organs ( $P < 0.05$ )



**Fig. 7** MBF1 mRNA expression in epidermis, fat body and middle gut of fifth instar. The levels of each MBF1 mRNA and actin were measured by SYBR Green I real-time RT-PCR

**Acknowledgements** This work was supported by the Hi-Tech Research and Development Program of China (No. 2008AA10Z132 and No. 2006AA10A119).

## References

- Johnson P, McKnight S (1989) Eukaryotic transcriptional regulatory proteins. *Annu Rev Biochem* 58:799–839. doi:10.1146/annurev.bi.58.070189.004055
- Mitchell PJ, Tjian R (1989) Transcriptional regulation in mammalian cells by sequence-specific DNA binding proteins. *Science* 245:371–378. doi:10.1126/science.2667136
- Lewin B (1990) Commitment and activation at pol II promoters: a tail of protein–protein interactions. *Cell* 61:1161–1164. doi:10.1016/0092-8674(90)90675-5
- Torchia J, Glass C, Rosenfeld MG (1998) Co-activators and co-repressors in the integration of transcriptional responses. *Curr Opin Cell Biol* 10:373–383. doi:10.1016/S0955-0674(98)80014-8
- Kingston RE (1999) A shared but complex bridge. *Science* 399:199–200
- Mannervik M, Nibu Y, Zhang H, Levine M (1999) Transcriptional coregulators in development. *Science* 284:606–609. doi:10.1126/science.284.5414.606
- Takemaru K, Harashima S, Ueda H, Hirose S (1998) Yeast coactivator MBF1 mediates GCN4-dependent transcriptional activation. *Mol Cell Biol* 18:4971–4976
- Zhu G, LaGier MJ, Hirose S, Keithly JS (2000) *Cryptosporidium parvum*: functional complementation of a parasite transcriptional coactivator CpMBF1 in yeast. *Exp Parasitol* 96:195–201. doi:10.1006/expr.2000.4574
- Ueda H, Sun GC, Murata T, Hirose S (1992) A novel DNA-binding motif abuts the zinc finger domain of insect nuclear hormone receptor FTZ-F1 and mouse embryonic long terminal repeat-binding protein. *Mol Cell Biol* 12(12):5667–5672
- Sun GC, Hirose S, Ueda H (1994) Intermittent expression of BmFTZ-F1, a member of the nuclear hormone receptor superfamily during development of the silkworm *Bombyx mori*. *Dev Biol* 162(2):426–437. doi:10.1006/dbio.1994.1099
- Goldsmith MR, Shimada T, Abe H (2005) The genetics and genomics of the silkworm, *Bombyx mori*. *Annu Rev Entomol* 50:71–100. doi:10.1146/annurev.ento.50.071803.130456
- Nagaraju J, Goldsmith MR (2002) Silkworm genomics-progress and prospects. *Curr Sci* 83(4):415–425
- Kiguchi K, Agui N (1981) Ecdysteroid levels and developmental events during larval molting in the silkworm, *Bombyx mori*. *J Insect Physiol* 27:805–812. doi:10.1016/0022-1910(81)90072-X
- Takemaru K, Li FQ, Ueda H, Hirose S (1997) Multiprotein bridging factor 1 (MBF1) is an evolutionarily conserved transcriptional coactivator that connects a regulatory factor and TATA element-binding protein. *Proc Natl Acad Sci USA* 94:7251–7256. doi:10.1073/pnas.94.14.7251
- Liu QX, Ueda H, Hirose S (2000) MBF2 is a tissue- and stage-specific coactivator that is regulated at the step of nuclear transport in the silkworm *Bombyx mori*. *Dev Biol* 225:437–446. doi:10.1006/dbio.2000.9836
- Kabe Y, Goto M, Shima D, Imai T, Wada T, Morohashi K et al (1999) The role of human MBF1 as a transcriptional coactivator. *J Biol Chem* 274:34196–34202. doi:10.1074/jbc.274.48.34196
- Gerace L (1995) Nuclear export signals and the fast track to the cytoplasm. *Cell* 82(3):341–344. doi:10.1016/0092-8674(95)90420-4
- Li FQ, Takemaru K, Goto M, Ueda H, Handa H, Hirose S (1997) Transcriptional activation through interaction of MBF2 with TFIIA. *Genes Cells* 2:143–153. doi:10.1046/j.1365-2443.1997.1090306.x
- Li FQ, Ueda H, Hirose S (1994) Mediators of activation of fushi tarazu gene transcription by BmFTZ-F1. *Mol Cell Biol* 14(5):3013–3021
- Liu QX, Jindra M, Ueda H, Hiromi Y, Hirose S (2003) *Drosophila* MBF1 is a co-activator for tracheae defective and contributes to the formation of tracheal and nervous systems. *Development* 130:719–728. doi:10.1242/dev.00297
- Tsuda K, Tsuji T, Hirose S, Yamazaki K (2004) Three Arabidopsis MBF1 homologs with distinct expression profiles play roles

- as transcriptional co-activators. *Plant Cell Physiol* 45:225–231. doi:[10.1093/pcp/pch017](https://doi.org/10.1093/pcp/pch017)
22. Ozaki J, Takemaru KI, Ikegami T, Mishima M, Ueda H, Hirose S et al (1999) Identification of the core domain and the secondary structure of the transcriptional coactivator MBF1. *Genes Cells* 4:415–424. doi:[10.1046/j.1365-2443.1999.00267.x](https://doi.org/10.1046/j.1365-2443.1999.00267.x)
23. Jindra M, Gaziova I, Uhlirova M, Okabe M, Hiromi Y, Hirose S (2004) Coactivator MBF1 preserves the redox-dependent AP-1 activity during oxidative stress in *Drosophila*. *EMBO J* 23:3538–3547. doi:[10.1038/sj.emboj.7600356](https://doi.org/10.1038/sj.emboj.7600356)
24. Brendel C, Gelman L, Auwerx J (2002) Multiprotein bridging factor-1 (MBF-1) is a cofactor for nuclear receptors that regulate lipid metabolism. *Mol Endocrinol* 16(6):1367–1377. doi:[10.1210/me.16.6.1367](https://doi.org/10.1210/me.16.6.1367)