



Research paper

Effects of zinc-exchanged montmorillonite with different zinc loading capacities on growth performance, intestinal microbiota, morphology and permeability in weaned piglets

L.F. Jiao, Y.L. Ke, K. Xiao, Z.H. Song, J.J. Lu^{*}, C.H. Hu^{*}

Animal Science College, Zhejiang University, The Key Laboratory of Molecular Animal Nutrition, Ministry of Education, Hangzhou 310058, China



ARTICLE INFO

Article history:

Received 12 November 2014

Received in revised form 17 March 2015

Accepted 14 April 2015

Available online 25 May 2015

Keywords:

Zinc-exchanged montmorillonite

Intestinal microbiota

Barrier integrity

Diarrhea

Weaned piglets

ABSTRACT

Three zinc-exchanged montmorillonites (Zn-Mt1, Zn-Mt2, and Zn-Mt3, the amount of Zn adsorbed on Mt was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively) were used to study the effects on growth performance, diarrhea, intestinal microbiota and barrier function in weaned piglets. A total of 144 piglets (Duroc $\times$ Landrace $\times$ Yorkshire), weaned at 21 ± 1 d with average initial body weight of 6.5 kg were allotted to four treatments groups for two weeks. Four treatments were: (1) control: basal diet; (2) ZM1: basal diet + 150 mg/kg Zn as Zn-Mt1; (3) ZM2: basal diet + 150 mg/kg Zn as Zn-Mt2; and (4) ZM3: basal diet + 150 mg/kg Zn as Zn-Mt3. Results showed that, compared with the control, supplementation of Zn-Mt2 or Zn-Mt3 improved ($P < 0.05$) average daily gain (ADG) and average daily feed intake (ADFI), and lowered ($P < 0.05$) fecal scores. Supplemental Zn-Mt2 or Zn-Mt3 increased ($P < 0.05$) villus height and the ratio of villus height and crypt depth at the jejunal mucosa. Supplementation of Zn-Mt2 or Zn-Mt3 decreased ($P < 0.05$) the number of *Escherichia coli* and *Streptococcus suis*, the paracellular permeability of fluorescein isothiocyanate dextran 4 kDa (FD4), but increased transepithelial electrical resistance (TER) in jejunum and colon of weaned piglets. No difference was observed between the Zn-Mt1 group and the control group. The results indicated that the in vivo efficacy of three Zn-Mts was different. Supplementation of Zn-Mt2 was as effective as Zn-Mt3 in enhancing growth performance, alleviating diarrhea, as well as improving intestinal microbiota, barrier function of weaned pigs under our trial conditions.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Weaning is the most significant event in the life of pigs and weaning process is commonly associated with microbial imbalance and intestinal barrier dysfunction, which are responsible for the stunted growth and diarrhea observed in the first 2 weeks after weaning (Moeser et al., 2007; Hu et al., 2013c; Xiao et al., 2014). Feeding pharmacological level of Zn (2000–4000 mg/kg) is used in the pig industry to alleviate post-weaning diarrhea and improve performance (Hu et al., 2014). However, the strategy has been criticized because high level of Zn results in large quantities of Zn excreted and poses an environmental problem (Carlson et al., 2004). Therefore, it is essential to find an alternative to reduce Zn supplementation for sustaining swine production.

Abbreviations: ADFI, average daily feed intake; ADG, average daily gain; SEM, standard error of the mean; Zn-Mt, zinc-exchanged montmorillonite; FD4, fluorescein isothiocyanate dextran 4 kDa; TER, transepithelial electrical resistance.

^{*} Corresponding authors.

E-mail addresses: jjlu@zju.edu.cn (J.J. Lu), chhu@zju.edu.cn (C.H. Hu).

Special attention has been paid to natural clay minerals, which are used as a potential carrier for release of active ingredients in controlled drug delivery systems (Rodrigues et al., 2013; Ke et al., 2014). A few researches have reported that zinc-bearing clinoptilolite may be used as an antibiotic growth promoter substitution for broiler chicken (Wang et al., 2012; Tang et al., 2014). Zinc-exchanged montmorillonite (Zn-Mt) is prepared by an ion-exchange reaction and studies are mainly focused on the physical and chemical properties in vitro (Malachová et al., 2011). However, there are no data regarding the biological effects of Zn-Mt in weaned pigs so far. Additionally, it has been reported that Zn-Mts with different Zn loading capacities had different physical and chemical properties. Churakov and Dähn (2012) found that whether Zn was adsorbed on montmorillonite (Mt) external surface or within its interlayer depended on the amount of Zn loading. Dähn et al. (2011) found that Zn-Mt with low or medium Zn loading capacity had different structures. Shi et al. (2010) reported that the Zn content in the interlayer space of Mt had an effect on the antibacterial activity of Zn-Mt. So we hypothesize that the different Zn loading capacities will affect in vivo efficiency.

The aim of the present study is to investigate the effects of Zn-Mt with different Zn loading capacities on growth performance, intestinal microbiota, morphology and permeability in weaned piglets.

2. Materials and methods

2.1. Materials

Mt was obtained from the Inner Mongolia Autonomous Region, China. The content of the purified Mt was 99.0%. Zn-Mt was prepared by Zn^{2+} -exchanged reaction. Ten grams of the Mt was mixed with 0.1 L of 0.2 mol/L NaCl solution. The dispersion was agitated for 5 h on a magnetic stirrer (700 rpm) at 25 °C. The Na-Mt was then separated by centrifugation at a speed of 8000 g for about 15 min and washed with deionized water for three times. The washed Na-Mt was then added to ZnSO_4 solution at 0.20 mol/L, 0.15 mol/L and 0.10 mol/L, respectively. The dispersion was agitated at 60 °C for 6 h on a magnetic stirrer (700 rpm). After centrifugation at 8000 g for 5 min, the sediment was washed with deionized water for three times, dried at 80 °C for 24 h, then ground to a size less than 50 μm pore diameter. Zn concentration in the Zn-Mt1, Zn-Mt2 and Zn-Mt3, as measured by atomic absorption spectroscopy (ICE 3300, Thermo Fisher Scientific, Waltham, USA), was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively.

2.2. Experimental design and sample collection

All procedures were approved by the Institutional Animal Care and Use Committee of Zhejiang University. A total of 144 piglets (Duroc $\times$ Landrace $\times$ Yorkshire), weaned at 21 ± 1 d with average initial body weight of 6.5 kg were allotted to four treatments. The four treatments were: (1) control: basal diet; (2) ZM1: basal diet + 150 mg/kg Zn as Zn-Mt1; (3) ZM2: basal diet + 150 mg/kg Zn as Zn-Mt2; and (4) ZM3: basal diet + 150 mg/kg Zn as Zn-Mt3. Each treatment had six pens of six piglets. Basal diets were formulated according to the National Research Council (1998) (Table 1). All pigs were given ad libitum access to mash feed and water for two weeks. Average daily gain (ADG), average daily feed intake (ADFI), and feed/gain

were measured. Post-weaning scour score was monitored for each pig according to Hu et al. (2012).

At 14 d after weaning, six piglets from each treatment were killed based on average diarrhea score. The gastrointestinal tract was quickly removed. Segments (1 cm) of the proximal jejunum were fixed in 10% formalin for morphology measurements. Adjacent jejunum and proximal colon were prepared for Ussing chamber studies. The intestinal contents from jejunum and proximal colon were collected, rapidly frozen in liquid nitrogen and stored at -80 °C for microbiota analysis.

2.3. Sample analysis

16S ribosomal RNA-based methods were used for the abundances of *Streptococcus suis* and *Escherichia coli* as described by Su et al. (2008). Briefly, total DNA was extracted from jejunal and colonic contents using a TIANamp Stool DNA Kit (Tiangen Biotech, Beijing, China) according to the manufacturer's instructions. The amount and quality of DNA were measured at 260 and 280 nm using ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). The primers of *S. suis* and *E. coli* were shown in Table 2. Real-time PCR was performed on 7500 real time PCR systems (Applied Biosystems, Foster City, USA) using Fast SYBR® Green Master Mix (Applied Biosystems, Foster City, USA). Each standard dilution and sample were assayed in triplicate in a 20 μL reaction mixture containing 10 μL of Fast SYBR Green Master Mix, 1 μL 25 pmol/ μL of each primer and 7 μL Nuclease-free water and 1 μL 50 ng/ μL DNA template. PCR amplification was performed with an initial denaturation step of 95 °C for 10 min, 40 cycles of 95 °C for 3 s, 60 °C for 1 min. For SYBR-Green® amplifications, a melting step was added to improve amplification specificity. Standard curves were generated as described by Li et al. (2009). The concentration of 16S rRNA gene abundance was plotted against the Cycle threshold value (CT value).

Three cross-sections for each jejunal sample were stained with hematoxylin and eosin using standard paraffin embedding procedures. Crypt depth and villus height were measured in at least ten well-oriented crypt-villus units using image analysis (Leica Imaging Systems Limited) and averaged for each sample.

Proximal jejunum and colon were stripped from the seromuscular layer in oxygenated Ringer's solution. Tissues were mounted in EasyMount Ussing chamber system (model VCC MC6, Physiologic Instruments, San Diego, CA, USA) as described previously (Hu et al., 2013a). Briefly, the clamps were connected to Acquire and Analyse software (Physiologic Instruments, San Diego, CA) for automatic data collection. After a 15-min equilibration period on Ussing chambers, transepithelial electrical resistance (TER) was recorded at 15-min intervals over a 1-h period. The epithelial barrier function was measured by the fluxes of fluorescein isothiocyanate dextran 4 kDa (FD4). The probe FD4 (Sigma-Aldrich, St. Louis, MO) was added to the mucosal side at the final concentration of 0.4 mg/mL. The samples were taken from the serosal side of tissues. The concentration of FD4 was measured by a fluorescence microplate reader (FLx800, Bio-Tek Instruments, Inc.).

2.4. Statistical analysis

One-way analysis of variance (ANOVA) was conducted using SPSS 9.0 statistical package (SPSS Inc., Chicago, IL). Differences among means were tested using Duncan's multiple range tests. Effects were considered significant at $P < 0.05$.

Table 1
Ingredient and composition of basal diets on an as-fed basis.

Ingredients, g/kg	
Maize	572.5
Soybean meal, crude protein 450 g/kg	257
Fish meal, crude protein 600 g/kg	50
Spray-dried plasma protein, crude protein 750 g/kg	25
Dried whey, crude protein 125 g/kg	45
Soybean oil	20
Dicalcium phosphate	11
Limestone	5
Sodium chloride	3
L-Lysine HCl, 780 g/kg	1
DL-Methionine, 990 g/kg	0.5
Vitamin–mineral premix ^a	10
Analyzed composition, g/kg	
Digestible energy ^b , MJ/kg	14.38
Moisture	85.7
Crude protein	224.1
Lysine	14.1
Methionine	3.8
Calcium	8.6
Total phosphorus	6.7
Zn (mg/kg)	127.3

^a Supplied per kilogram of diet: vitamin A, 6000 IU; vitamin D₃, 600 IU; vitamin E, 50 IU; vitamin K₃, 1.5 mg; riboflavin, 8.0 mg; thiamine, 2.0 mg; niacin, 30 mg; pyridoxine, 3.0 mg; pantothenic acid, 20 mg; choline, 800 mg; folic acid, 0.6 mg; biotin, 0.10 mg; vitamin B₁₂, 0.04 mg; Zn (ZnSO_4), 100 mg; Cu ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$), 16 mg; I (KI), 0.2 mg; Fe (FeSO_4), 125 mg; Mn ($\text{MnSO}_4 \cdot \text{H}_2\text{O}$), 15 mg; Se (Na_2SeO_3), 0.3 mg.

^b Digestible energy was calculated from data provide by Feed Database in China (2012).

Table 2
PCR primers for the quantification of intestinal microbiota by real-time PCR.

Target organism	Forward/reverse	Sequence 5'–3'
<i>E. coli</i>	F	CATGCCCGCTGTATGAAGAA
	R	CGGTAACGTCAATGAGCAAA
<i>Streptococcus suis</i>	F	CAGTATTATCCGCATGGTAGATAT
	R	GTAAGATACCGTCAAGTGAGAA

Table 3

Effects of zinc-exchanged montmorillonite on growth performance and diarrhea index of weaned piglets.

Items	Control	ZM1 [*]	ZM2 [*]	ZM3 [*]	SEM ¹	P
Initial weight, kg	6.52	6.61	6.57	6.48	1.03	0.924
Final weight, kg	10.46	10.74	10.93	10.96	1.25	0.867
ADG ² , g/d	282 ^b	295 ^{ab}	312 ^a	315 ^a	9.06	0.035
ADFI ³ , g/d	362 ^b	379 ^{ab}	386 ^a	390 ^a	8.30	0.002
Feed/gain ratio	1.28	1.28	1.25	1.24	0.02	0.822
Fecal scores	3.17 ^a	2.50 ^b	1.83 ^c	1.50 ^c	0.20	0.012

^{a–b}Means within a row with different letters differ significantly ($P < 0.05$).^{*} Zinc-exchanged montmorillonite (Zn-Mt), Zn-Mt1, Zn-Mt2, and Zn-Mt3, the amount of Zn adsorbed on Mt was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively.¹ Standard error of means, $n = 6$.² ADG = Average daily gain.³ ADFI = Average daily feed intake.

3. Results

3.1. Growth performance and postweaning scour scores

Table 3 showed the growth performance and diarrhea index of weaned piglets. As compared with the control, supplementation with Zn-Mt2 or Zn-Mt3 increased ($P < 0.05$) ADG and ADFI, and decreased ($P < 0.05$) fecal scores. No significant difference ($P > 0.05$) was observed in the growth performance and scour scores between the Zn-Mt2 group and Zn-Mt3 group. The growth performance and scour scores of weaned pigs fed with Zn-Mt1 did not differ from those fed with the control diet. The feed/gain was not affected by dietary treatments ($P > 0.05$).

3.2. Intestinal microbiota

Supplementation with Zn-Mt2 or Zn-Mt3 reduced ($P < 0.05$) the number of *S. suis* and *E. coli* in the jejunum and proximal colon of weaned piglets as compared with the control (Table 4). No significant difference ($P > 0.05$) was observed between the Zn-Mt2 group and Zn-Mt3 group. The intestinal microbiota of weaned pigs fed with Zn-Mt1 did not differ from those fed with the control diet.

3.3. Intestinal morphology

Table 5 showed the jejunal morphology of weaned piglets. As compared with the control, supplementation with Zn-Mt2 or Zn-Mt3 increased villus height and the ratio of villus height and crypt depth ($P < 0.05$). No significant difference ($P > 0.05$) was observed between the Zn-Mt2 group and Zn-Mt3 group. The jejunal morphology of pigs fed Zn-Mt1 did not differ from those fed the control diet ($P > 0.05$).

3.4. Intestinal barrier function

Intestinal barrier function of weaned pigs, as reflected by the TER and paracellular flux of FD4, was presented in Table 6. As compared with the control, pigs fed Zn-Mt2 or Zn-Mt3 had a higher TER value

Table 4Effect of zinc-exchanged montmorillonite on intestinal microbiota of weaned piglets¹.

Items	Control	ZM1 [*]	ZM2 [*]	ZM3 [*]	SEM ²	P
Jejunum						
<i>E. coli</i>	7.04 ^a	6.67 ^{ab}	6.45 ^b	6.38 ^b	0.15	0.025
<i>Streptococcus suis</i>	6.14 ^a	5.78 ^{ab}	5.49 ^b	5.30 ^b	0.20	0.037
Colon						
<i>E. coli</i>	8.32 ^a	7.94 ^{ab}	7.53 ^b	7.39 ^b	0.22	0.034
<i>Streptococcus suis</i>	7.47 ^a	7.23 ^{ab}	6.71 ^b	6.56 ^b	0.22	0.029

^{a–b}Means within a row with different letters differ significantly ($P < 0.05$).^{*} Zinc-exchanged montmorillonite (Zn-Mt), Zn-Mt1, Zn-Mt2, and Zn-Mt3, the amount of Zn adsorbed on Mt was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively.¹ Bacterial numbers are expressed as log₁₀ (16S rRNA gene copies g⁻¹ wet weight).² Standard error of the mean, $n = 6$.**Table 5**

Effects of zinc-exchanged montmorillonite on jejunal morphology of weaned piglets.

Items	Control	ZM1 [*]	ZM2 [*]	ZM3 [*]	SEM ¹	P
Villus height, μm	729 ^b	754 ^{ab}	779 ^a	820 ^a	19.6	0.022
Crypt depth, μm	375	361	350	347	12.1	0.377
Villus height: crypt depth	1.95 ^b	2.09 ^b	2.23 ^a	2.36 ^a	0.048	<0.001

^{a–b}Means within a row with different letters differ significantly ($P < 0.05$).^{*} Zinc-exchanged montmorillonite (Zn-Mt), Zn-Mt1, Zn-Mt2, and Zn-Mt3, the amount of Zn adsorbed on Mt was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively.¹ Standard error of the mean, $n = 6$.

($P < 0.05$) and lower FD4 permeability in jejunum and colon ($P < 0.05$). No significant difference ($P > 0.05$) was observed between the Zn-Mt2 group and Zn-Mt3 group. The intestinal barrier function of weaned pigs fed with Zn-Mt1 did not differ from those fed the control diet ($P > 0.05$).

4. Discussion

In the present study, we found that providing 150 mg/kg Zn, supplementation of Zn-Mt2 or Zn-Mt3 alleviated postweaning diarrhea and improved growth performance in the weaned pigs. Additionally, feeding three Zn-Mts in diets gave variable results on the growth performance of weaned piglets, which indicated that the in vivo efficiency of the Zn-Mt with different Zn loading capacities was different. Preliminary experiments showed that weaned pigs fed Mt (the amount is equivalent to the Mt in the Zn-Mt1, Zn-Mt2, Zn-Mt3 treatment, respectively) had no significant effect on growth performance (data not shown). Our previous study had reported that supplementation with 500 mg/kg Zn from montmorillonite–ZnO hybrid or 600 mg/kg Zn from zeolite–ZnO hybrid could enhance growth performance and alleviate diarrhea of weaned pigs. The amount of Zn as ZnO adsorbed on Mt and zeolite is 25% and 28%, respectively (Hu et al., 2012, 2013b). It was suggested that Mt with lower Zn loading capacity may perform better in modifying the rate, time, or site of Zn release within the gastrointestinal tract and increased access of Zn to the intestinal tract, which would improve its biological roles in the hindgut and therefore benefit overall gut health and growth of animals.

Numerous investigations have found the imbalance of the intestinal microbiota in weaning piglets, which results in overgrowing pathogenic bacteria (*E. coli*, *S. suis*) and causes intestinal disorders (Castillo et al., 2007; Su et al., 2008; Xu et al., 2014). Previous researches have reported that Mt could be used as a carrier for inorganic antibacterial agents, such as Ag⁺, Cu²⁺ and Zn²⁺, which are loaded into Mt by ion exchange (Magana et al., 2008). The antibacterial properties of metallic ion-exchanged Mt on *E. coli* have been proved in vitro (Hu and Xia, 2006; Malachová et al., 2011). The current study demonstrated for the first time that dietary Zn-Mt2 or Zn-Mt3 supplementation decreased the population of *E. coli* and *S. suis* in intestinal contents of weaned pigs. Two possible mechanisms for the antibacterial activity were proposed. One model involved the adsorption of the bacteria and immobilization

Table 6

Effect of zinc-exchanged montmorillonite on intestinal barrier integrity of weaned pigs.

Items	Control	ZM1 [*]	ZM2 [*]	ZM3 [*]	SEM ³	P
Jejunum						
TER ¹ , $\Omega \text{ cm}^2$	50.17 ^b	53.17 ^{ab}	58.67 ^a	59.83 ^a	2.40	0.030
FD4 ² flux, $\mu\text{g cm}^{-2} \text{ h}^{-1}$	2.58 ^a	2.17 ^{ab}	1.68 ^b	1.65 ^b	0.25	0.046
Colon						
TER ¹ , $\Omega \cdot \text{cm}^2$	65.00 ^b	69.17 ^{ab}	73.17 ^a	75.33 ^a	2.49	0.040
FD4 ² flux, $\mu\text{g cm}^{-2} \text{ h}^{-1}$	2.18 ^a	1.74 ^{ab}	1.39 ^b	1.35 ^b	0.22	0.047

^{a–b}Means within a row with different letters differ significantly ($P < 0.05$).^{*} Zinc-exchanged montmorillonite (Zn-Mt), Zn-Mt1, Zn-Mt2, and Zn-Mt3, the amount of Zn adsorbed on Mt was 6.3×10^4 , 5.1×10^4 and 4.2×10^4 mg/kg, respectively.¹ TER = transepithelial electrical resistance.² FD4 = fluorescein isothiocyanate dextran 4 kDa.³ Standard error of means, $n = 6$.

on the surface of the Zn-Mt. Alternatively, Zn was released from the Mt structure and directly exerted its antimicrobial effect on the bacteria. Therefore, the antibacterial effect of Zn-Mt may be explained by interactions of Mt (Magana et al., 2008).

There is evidence to suggest that a weaning transition of piglets is commonly associated with quick changes in the intestinal mucosa (Ke et al., 2014). A shortening of the villus decreases the surface area for nutrient absorption. The crypt is the area where stem cells divide to permit the renewal of the villus, and a large crypt indicates fast tissue turnover and a high demand for new tissue (Hu et al., 2013a). Changes in intestinal morphology, such as shorter villi and deeper crypts, can lead to poor nutrient absorption, increased secretion in the gut, diarrhea, reduced disease resistance, and impaired overall performance (Jiao et al., 2014). In the present study, increases in villus height and villus height: crypt depth ratio were observed in the small intestinal mucosa of weaned pigs fed the diet supplemented with Zn-Mt2 or Zn-Mt3. Similarly, Tang et al. (2014) reported that zinc-bearing clinoptilolite modulated intestinal morphology in broiler chickens. In our present study, Zn-Mt2 or Zn-Mt3 may partially improve the intestinal morphology via lowering the number of *E. coli* and *S. suis*.

An intact intestinal barrier plays a central role in preventing the translocation of intestinal bacteria, and the entering of toxic or allergenic substances from the gut into the body (Wijten et al., 2011). Intestinal barrier function can be commonly assessed by many indices such as TER and the paracellular permeability of FD₄ (Jiao et al., 2014). It was reported that small-intestinal barrier function of pigs deteriorated after weaning (Smith et al., 2010; Wijten et al., 2011). In the present study, consistent with improved intestinal morphology, increased TER and decreased FD₄ flux were observed in the small intestinal mucosa of weaned pigs fed the diet supplemented with Zn-Mt2 or Zn-Mt3. Zn is known to play an important role in maintaining barrier function of the gastrointestinal tract (Hu et al., 2014). Mt has high surface area and stand-out adhesive ability. These inherent advantages make Mt effectively act by attaching Zn²⁺ to the mucus to reinforce the intestinal mucosal barrier and help in the regeneration of the epithelium (Hu et al., 2012).

5. Conclusion

Providing 150 mg/kg Zn, the in vivo efficacy of Zn-Mt1, Zn-Mt2, and Zn-Mt3 was different in weaned piglets. Supplementation of Zn-Mt2 with 5.1×10^4 mg/kg Zn loading capacity or Zn-Mt3 with 4.2×10^4 mg/kg Zn loading capacity increased growth performance, alleviated postweaning diarrhea, and improved intestinal microflora and barrier function of weanling pigs. These positive effects of Zn-Mt may be attributed to Zn and montmorillonite interactions.

Acknowledgment

This work was jointly supported by the Special Fund for Agro-scientific Research in the Public Interest (No. 201403047).

References

- Carlson, M.S., Boren, C.A., Wu, C., Huntington, C.E., Bollinger, D.W., Veum, T.L., 2004. Evaluation of various inclusion rates of organic zinc either as apolysaccharide or proteinate complex on the growth performance, plasma, and excretion of nursery pigs. *J. Anim. Sci.* 82, 1359–1366.
- Castillo, M., Martín-Orúe, S.M., Nofrías, M., Manzanilla, E.G., Gasa, J., 2007. Changes in caecal microbiota and mucosal morphology of weaned pigs. *Vet. Microbiol.* 124, 239–247.
- Churakov, S.V., Dähn, R., 2012. Zinc adsorption on clays inferred from atomistic simulations and EXAFS spectroscopy. *Environ. Sci. Technol.* 46, 5713–5719.
- Dähn, R., Bradbury, M., Baeyens, B., 2011. Investigation of the different binding edge sites for Zn on montmorillonite using P-EXAFS—the strong/weak site concept in the 2SPNE SC/CE sorption model. *Geochim. Cosmochim. Acta* 75, 5154–5168.
- Feed Database in China, 2012. Table of Feed Composition and Nutritive Value in China. Twentieth ed. China Feed, Beijing, China.
- Hu, C.H., Xia, M.S., 2006. Adsorption and antibacterial effect of copper-exchanged montmorillonite on *Escherichia coli* K88. *Appl. Clay Sci.* 31, 180–184.
- Hu, C.H., Gu, L.Y., Luan, Z.S., Song, J., Zhu, K., 2012. Effects of montmorillonite–zinc oxide hybrid on performance, diarrhea, intestinal permeability and morphology of weanling pigs. *Anim. Feed Sci. Technol.* 177, 108–115.
- Hu, C.H., Song, J., Li, Y.L., Luan, Z.S., Zhu, K., 2013a. Diosmectite–zinc oxide composite improves intestinal barrier function, modulates expression of pro-inflammatory cytokines and tight junction protein in early weaned pigs. *Br. J. Nutr.* 110, 681–688.
- Hu, C.H., Xiao, K., Song, J., Luan, Z.S., 2013b. Effects of zinc oxide supported on zeolite on growth performance, intestinal microflora and permeability, and cytokines expression of weaned pigs. *Anim. Feed Sci. Technol.* 181, 65–71.
- Hu, C.H., Xiao, K., Luan, Z.S., Song, J., 2013c. Early weaning increases intestinal permeability, alters expression of cytokine and tight junction protein, and activates mitogen-activated protein kinases in pigs. *J. Anim. Sci.* 91, 1094–1101.
- Hu, C.H., Song, Z.H., Xiao, K., Song, J., Jiao, L.F., Ke, Y.L., 2014. Zinc oxide influences intestinal integrity, the expressions of genes associated with inflammation, and TLR4-myeloid differentiation factor 88 signaling pathways in weanling pigs. *Innate Immun.* 20, 478–486.
- Jiao, L.F., Song, Z.H., Ke, Y.L., Xiao, K., Hu, C.H., Shi, B., 2014. Cello-oligosaccharide influences intestinal microflora, mucosal architecture and nutrient transport in weaned pigs. *Anim. Feed Sci. Technol.* 195, 85–91.
- Ke, Y.L., Jiao, L.F., Song, Z.H., Xiao, K., Lai, T.M., Lu, J.J., Hu, C.H., 2014. Effects of cetylpyridinium-montmorillonite, as alternative to antibiotic, on the growth performance, intestinal microflora and mucosal architecture of weaned pigs. *Anim. Feed Sci. Technol.* 198, 257–262.
- Li, M., Penner, G.B., Hernandez-Sanabria, E., Oba, M., Guan, L.L., 2009. Effects of sampling location and time, and host animal on assessment of bacterial diversity and fermentation parameters in the bovine rumen. *J. Appl. Microbiol.* 107, 924–934.
- Magana, S.M., Quintana, P., Aguilar, D.H., Toledo, J.A., Angeles-Chavez, C., Cortes, M.A., Leon, L., Freile-Pelegrin, Y., Lopez, T., Torres Sanchez, R.M., 2008. Antibacterial activity of montmorillonites modified with silver. *J. Mol. Catal. A Chem.* 281, 192–199.
- Malachová, K., Praus, P., Rybková, Z., Kozák, O., 2011. Antibacterial and antifungal activities of silver, copper and zinc montmorillonites. *Appl. Clay Sci.* 53, 642–645.
- Moesser, A.J., Ryan, K.A., Nighot, P.K., Blikslager, A.T., 2007. Gastrointestinal dysfunction induced by early weaning is attenuated by delayed weaning and mast cell blockade in pigs. *Am. J. Physiol. Gastrointest. Liver* 293, G413–G421.
- National Research Council, 1998. Nutrient Requirements of swine. 10th Ed. National Academy Press, Washington, DC.
- Rodrigues, L.A.D.S., Figueiras, A., Veiga, F., de Freitas, R.M., Nunes, L.C., da Silva Filho, E.C., da Silva Leite, C.M., 2013. The systems containing clays and clay minerals from modified drug release: a review. *Colloids Surf. B* 103, 642–651 (COLLOID SURFACE B).
- Shi, Q.S., Tan, S.Z., Yang, Q.H., Jiao, Z.P., Ouyang, Y.S., Chen, Y.B., 2010. Preparation and characterization of antibacterial Zn²⁺-exchanged montmorillonites. *J. Wuhan Univ. Technol.* 25, 725–729.
- Smith, F., Clark, J.E., Overman, B.L., Tozel, C.C., Huang, J.H., Rivier, J.E., Blikslager, A.T., Moesser, A.J., 2010. Early weaning stress impairs development of mucosal barrier function in the porcine intestine. *Am. J. Physiol. Gastrointest. Liver* 298, G352–G363.
- Su, Y., Yao, W., Perez-Gutierrez, O.N., Smidt, H., Zhu, W.Y., 2008. 16S ribosomal RNA-based methods to monitor changes in the hindgut bacterial community of piglets after oral administration of *Lactobacillus sobrius* S1. *Anaerobe* 14, 78–86.
- Tang, Z.G., Wen, C., Wang, L.C., Wang, T., Zhou, Y.M., 2014. Effects of zinc-bearing clinoptilolite on growth performance, cecal microflora and intestinal mucosal function of broiler chickens. *Anim. Feed Sci. Technol.* 189, 98–106.
- Wang, L.C., Zhang, T.T., Wen, C., Jiang, Z.Y., Wang, T., Zhou, Y.M., 2012. Protective effects of zinc-bearing clinoptilolite on broilers challenged with *Salmonella pullorum*. *Poult. Sci.* 91, 1838–1845.
- Wijten, P.J., van der Meulen, J., Verstegen, M.W., 2011. Intestinal barrier function and absorption in pigs after weaning: a review. *Br. J. Nutr.* 105, 967–981.
- Xiao, K., Song, Z.H., Jiao, L.F., Ke, Y.L., Hu, C.H., 2014. Developmental changes of TGF- β 1 and Smads signaling pathway in intestinal adaption of weaned pigs. *PLoS ONE* 9 (e104589).
- Xu, J.X., Xu, C.C., Chen, X.L., Cai, X., Yang, S.F., Sheng, Y.S., Wang, T., 2014. Regulation of an antioxidant blend on intestinal redox status and major microbiota in early weaned piglets. *Nutrition* 30, 584–589.