



## Mechanism of anti-angiogenic activities of chitooligosaccharides may be through inhibiting heparanase activity

Haizhi Quan<sup>a</sup>, Fei Zhu<sup>a</sup>, Xinyan Han<sup>a</sup>, Zirong Xu<sup>a,\*</sup>, Yilong Zhao<sup>b</sup>, Zhiguo Miao<sup>c</sup>

<sup>a</sup>Key Laboratory for Molecular Animal Nutrition of Ministry of Education, Feed Animal Sciences Institute, Zhejiang University, Qiutao North Road 164, Zhejiang, Hangzhou 310029, China

<sup>b</sup>Veterinarian Department Agriculture College Yanbian University, Longjing 133400, China

<sup>c</sup>Henan Institute of Science and Technology, College of Animal Science, Xinxiang, China

### ARTICLE INFO

#### Article history:

Received 25 February 2009

Accepted 28 February 2009

### SUMMARY

Metastatic disease is the primary cause of death for most cancer patients. Angiogenesis is the formation of a new capillary network from pre-existing vessels and required for tumor vasculature. Heparanase, a  $\beta$ -endoglucuronidase, assists tumor invasion, metastasis and angiogenesis. Chitooligosaccharides (COS) is obtained by hydrolysis of chitosan. COS has been proved to be anti-angiogenesis activity. The mechanism of COS inhibits angiogenesis is not very clear, COS is hypothesized by author to be an inhibitor of heparanase.

© 2009 Elsevier Ltd. All rights reserved.

### Introduction

Angiogenesis is the formation of a new capillary network from pre-existing vessels and is required for normal and tumor vasculature [1]. During angiogenesis upon stimulation by angiogenic growth factors and cytokines, quiescent vascular endothelial cells degrade the underlying basement membrane, migrate into the extracellular matrix, proliferate, and form capillary structures [2]. Therefore, during tumor growth and metastasis, angiogenesis is required for proper nourishment, removal of metabolic wastes from tumor sites, and creating vessels for tumor cells migration [3,4].

Heparan sulphate proteoglycans (HSPGs), a key component of basement (BM) and extracellular matrix (ECM), is prominent components of blood vessels [5]. HSPGs themselves act as a storage depot for a number of cytokines and growth factors such as basic fibroblast growth factor (bFGF), vascular endothelial growth factor (VEGF). HSPGs also bind specifically to the heparan sulphate (HS) glycosaminoglycan (GAG) chains [6]. Heparanase is a  $\beta$ -endoglucuronidase, which acts via a hydrolytic mechanism to cleave glycosidic bonds in HS [7]. A critical early event in the tumor angiogenic process is that heparanase degrades the HS of HSPG in BM and ECM, followed by endothelial cells migration toward the angiogenic stimulus. Moreover by releasing HS-bound angiogenic growth factors (i.e. VEGF) from the ECM, heparanase may indirectly facilitate endothelial cells migration and proliferation. The processes heparanase assists tumor invasion and metastasis are just like those of endothelial cells migration and proliferation

[8]. So heparanase plays an important role in tumor invasion, metastasis and angiogenesis.

Chitosan is the deacetylated form of chitin, and is a linear polymer of *N*-acetyl- $\beta$ -glucosamine and deacetylated glucosamine. The molecular structure of chitosan makes it shares some characteristics of GAG [9]. Chitooligosaccharides (COS) can be obtained by either chemical or enzymatic hydrolysis of chitosan. COS are composed of 3–10 *N*-acetyl-glucosamine (NAGA) or glucosamine residues [10]. In biomedicine field, COS is more applicable than chitosan because of its water solubility and favorable properties of biodegradability, biocompatibility and other bioactivities [11]. COS has been proved some characteristics different from chitosan such as anti-angiogenesis [12], inhibiting vascular endothelium cells migration [13], depressing tumor invasion [14].

COS is a successful inhibitor of tumor induced angiogenesis [12]. Nevertheless the mechanism of COS anti-angiogenesis is not very clear.

### Hypothesis

COS may be a competitive inhibitor for heparanase, due to its molecular structure similar to that of HS GAG, the substrate of heparanase.

### Evaluation of hypothesis

Metastatic disease is the primary cause of death for most cancer patients [15]. Tumor growth and metastasis require angiogenesis when the tumor reaches 1–2 mm in diameter [16]. Heparanase activity was implicated in cellular invasion associated with tumor angiogenesis and metastasis [17]. Downregulating the expression

\* Corresponding author. Tel.: +86 571 86971075.

E-mail address: [zrxu@zju.edu.cn](mailto:zrxu@zju.edu.cn) (Z. Xu).

of heparanase inhibits the invasion, angiogenesis and metastasis of human hepatocellular carcinoma [18]. PI-88, a heparanase inhibitor significantly reduces the expression of VEGF receptor A [19]. Excess heparanase expression is a poor prognostic indicator of breast and gastric cancer [20,21]. It can be concluded that heparanase inhibitor inhibits tumor growth, metastasis and angiogenesis. Therefore heparanase inhibitor is a direction for drug development in cancer therapy [6].

Low molecular weight heparin has been used to control angiogenesis and heparanase activity in anti-tumor drug discovery [22,23]. However chitin, which is obtained in large quantities from crustacean shells, the material of COS, has abundant resource in nature [24]. Maybe it is a potent predominance of COS drug discovery. Besides, COS can be a safely oral administered drug because of its few side effects [25], water solubility [11] and epithelial cells permeability [26].

## References

- [1] Dass CR, Tran TR, Choong PFM. Angiogenesis inhibitor and the need for anti-angiogenic therapeutics. *Crit Rev Oral Biol Med* 2007;10:927–36.
- [2] Toshikazu N, Kunio M. Angiogenesis inhibitors: from laboratory to clinical application. *Biochim Biophys Res Commun* 2005;333:289–91.
- [3] Tiziana T, Francesca R, Pier PC. Molecular basis of angiogenesis and cancer. *Oncogene* 2003;22:6549–56.
- [4] Douglas H, Judah F. Patterns and emerging mechanisms of the angiogenic switch during tumorigenesis. *Cell* 1996;86:353–64.
- [5] Rohloff J, Zinke J, Schoppmeyer K, et al. Heparanase expression is a prognostic indicator for postoperative survival in pancreatic adenocarcinoma. *Br J Cancer* 2002;86:1270–5.
- [6] McKenzied EA. Heparanase: a target for drug discovery in cancer and inflammation. *Br J Pharmacol* 2007;151:1–14.
- [7] Martin G, George WY. Heparanase, hyaluronan, and CD44 in cancers: a breast carcinoma perspective. *Cancer Res* 2006;21:10233–7.
- [8] Neta I, Michael E, Israel V. Regulation, function and clinical significance of heparanase in cancer metastasis and angiogenesis. *Int J Biochem Cell Biol* 2006;38:2018–39.
- [9] Sur JK, Matthew HW. Application of chitosan-based polysaccharide biomaterials in cartilage tissue engineering: a review. *Biomaterial* 2000;21:2589–98.
- [10] Yanping H, Luhang Z, Zhijun Y, Jie F, Qiqi Y. Role of mannose receptor in oligochitosan-mediated stimulation of macrophage function. *Int Immunopharmacol* 2005;5:1533–42.
- [11] Haige W, Ziang Y, Xuefang B, Yuguang D, Bingcheng L. Anti-angiogenic activities of chitooligosaccharides. *Carbohydr Polym* 2008;73:105–10.
- [12] Harish Prashanth KV, Tharanathan RN. Depolymerized products of chitosan as potent inhibitors of tumor-induced angiogenesis. *Biochim Biophys Acta* 2005;1722:22–9.
- [13] Yoshiharu O, Midori W, Katsuyuki M, et al. Effects of chitin/chitosan and their oligomers/monomers on migrations of fibroblasts and vascular endothelium. *Biomaterial* 2002;23:1975–9.
- [14] Moon Moo K, Se Kwon K. Chitooligosaccharides inhibit activation and expression of matrix metalloproteinase-2 in human dermal fibroblasts. *FEBS Lett* 2006;580:2261–6.
- [15] Patricia SS. Tumor metastasis: mechanistic insights and clinical challenges. *Nat Med* 2006;12:895–904.
- [16] Binghua J, Lingzhi L. PI3K/PTEN signaling in tumorigenesis and angiogenesis. *Biochim Biophys Acta* 2008;1784:150–8.
- [17] Motowo N, Tatsuro I, Daniela DF, Nicola DF, Garth LN. Heparan sulfate degradation: relation to tumor invasive and metastatic properties of mouse B16 melanoma sublines. *Science* 1983;220:611–3.
- [18] Youlei Z, Li L, Yi W, et al. Downregulating the expression of heparanase inhibits the invasion, angiogenesis and metastasis of human hepatocellular carcinoma. *Biochem Biophys Res Commun* 2007;358:124–9.
- [19] Johanna AJ, Craig F, Nicole MM, et al. *Oncogene* 2002;24:4037–51.
- [20] Justin B, Maxhimer BA, Roderick M, et al. Heparanase-1 expression is associated with the metastatic potential of breast cancer. *Surgery* 2002;132:326–33.
- [21] Munenori T, Yoshio N, Takaomoto O, et al. Heparanase expression correlates with invasion and poor prognosis in gastric cancers. *Lab Invest* 2003;83:613–22.
- [22] Judah F, Paul BW, Madeleine MJ. Control of angiogenesis with synthetic heparin substitutes. *Science* 1989;243:1490–3.
- [23] Rebecca L, Clive PP. Novel drug development opportunities for heparin. *Nat Rev Drug Dis* 2002;1:140–8.
- [24] Hyun Joong Y, Myoung EM, Haeng Soon P, Suh Young I, Young Ho K. Chitosan oligosaccharide (COS) inhibits LPS-induced inflammatory effects in RAW264.7 macrophage cells. *Biochem Biophys Res Commun* 2007;358:954–9.
- [25] Vale'rie D, Amin KM, June RM. Effect of chitosan on epithelial permeability and structure. *Int J Pharm* 1999;182:21–32.
- [26] Jung Kul L, Hyun Soo L, Jung Hoe K. Cytotoxic activity of aminoderivatized cationic chitosan derivatives. *Bioorg Med Chem Lett* 2002;12:2949–51.