

ORIGINAL ARTICLE

Effect of the extract made from *Cochinchina momordica* seeds on the humoral immune responses of mice to a commercial foot-and-mouth disease vaccine (serotypes O and Asia 1)

Kedsirin Sakwiwatkul^{1,3}, Rui-li Li¹, Xiao-ming Song¹ and Song-hua Hu^{1,2}

¹Department of Veterinary Medicine, College of Animal Sciences, Zhejiang University, Hangzhou, Zhejiang 310029, China ²Key Laboratory of Animal Epidemic Etiology & Immunological Prevention of Ministry of Agriculture, P. R. China, Hangzhou, Zhejiang 310029, China ³Faculty of Veterinary Medicine and Animal Science, Mahasarakham University, Talad, Mahasarakham 44000, Thailand

ABSTRACT

The extract from ECMS was investigated for its effect on the humoral immune responses to foot-and-mouth disease vaccination. Fifty-six mice were randomly divided into seven groups with eight animals in each. Mice in groups 5 to 7 were subcutaneously (s.c.) injected with 0.5 mg DEX daily for 4 days to induce immunosuppression. The animals were then orally given ECMS (200 µg in 250 µl saline) in groups 3 and 6 or 250 µl saline in group 2, or s.c. injected with ECMS (50 µg in 100 µl saline) in groups 4 and 7 or 100 µl saline in group 5. After that, the animals in groups 2 to 7 were s.c. immunized twice with 100 µl of commercial oil-adjuvanted bivalent FMDV vaccine (serotypes O and Asia 1) at intervals of 21 days. Mice in group 1 received injection of 100 µl saline only. After 2 weeks, blood was sampled to determine FMDV-specific IgG and isotype IgG1, IgG2a, IgG2b and IgG3. Results indicated that oral administration or s.c. injection of ECMS augmented responses of specific IgG and most IgG isotypes. Giving ECMS tended to enhance serum-specific IgG and IgG isotype responses of mice immunosuppressed by s.c. injection of DEX. Considering the safety and immunomodulatory effect of ECMS in both normal and immunosuppressed mice demonstrated in the present study, this extract deserves further investigation to evaluate its potential in improving FMD vaccination in farm animals such as pigs, sheep and cattle.

Key words *Cochinchina momordica* seeds, foot-and-mouth disease, vaccine.

Foot-and-mouth disease is caused by FMDV, which is the prototype member of the *Aphthovirus* genus, *Picornaviridae* (1) and occurs as seven major serotypes: A, O, C, Asia 1, SAT 1, SAT 2 and SAT 3, but a large number of subtypes are found within each serotype (2, 3). The disease mainly affects the cloven-hoofed animals. FMD has been recorded as one of the most important animal diseases in the world by the World Organization for Animal Health (OIE) due to the rapid transmission of FMDV among susceptible

animals (4). Vaccination is a common practice against FMD in many countries. However, failure to elicit effective immune responses by vaccination has been frequently reported (5–7). In agreement with this, Hao *et al.* (8) observed that only 31.9% out of 91 pigs vaccinated against FMD (type O) produced serum antibody titers needed for immune protection. Hence, currently available vaccines should be improved in order to effectively prevent infectious diseases in animals. Higher dosage of vaccine

Correspondence

Song-hua Hu, 268 Kaixuan Rd, Hangzhou, Zhejiang 310029, P. R. China.
Tel: 0086 0571 8697 1852; fax: 0086 0571 8697 1852; email: songhua@zju.edu.cn

Received 10 August 2009; revised 17 September 2009; accepted 23 September 2009.

List of Abbreviations: DEX, dexamethasone; ECMS, *Cochinchina momordica* seeds; FMD, foot-and-mouth disease; FMDV, FMD virus; Ig, immunoglobulin; OVA, ovalbumin.

or higher frequency of vaccination and use of adjuvants such as saponins and a variety of immunomodulators are some approaches to improve the immune response to vaccination (9–14).

Semen momordicae is the seed of *Momordica cochinchinensis* (Lour.) Spreng. mainly growing in Southeast Asian countries and southern China (15). Traditionally, the seeds are used to treat inflammatory swelling, diarrhea and suppurative skin infections in human beings and animals (16). From the seeds, Iwamoto *et al.* (17) have isolated momordica saponins I and II. According to the chemical analysis, momordica saponin I is a triterpenoid saponin containing disaccharide chains, and momordica saponin II is structurally similar to quillaic acid. Our previous study has shown that the addition of (ECMS extract in a commercial foot-and-mouth disease vaccine can significantly enhance immune responses in pigs (18). ECMS given s.c. has also been shown to exert an adjuvant effect on the immune response particularly at a humoral level to OVA (19) in mice and to influenza vaccination (H5N1) in chickens (20). In the present study, the immunomodulatory effects of ECMS by the oral or the s.c. route on the humoral immune response of normal and immunosuppressive mice against vaccination of bivalent FMDV serotypes O and Asia I vaccine were evaluated. To induce immunosuppression in ICR mice, DEX was injected s.c. prior to giving ECMS and bivalent FMDV vaccine, as DEX has been shown to inhibit several aspects of cell-mediated immunity, including antigen- and mitogen-induced lymphocyte proliferation (21) and is commonly used to produce an immunosuppressive model in mice.

MATERIALS AND METHODS

Experimental animals

Female ICR mice with a bodyweight of 19–22 g were purchased from Shanghai Laboratory Animal Center Co. Ltd (Shanghai, China) and kept in cages bedded with sawdust

in a controlled environment. Feed and water were supplied *ad libitum*. The study was conducted in accordance with the Guidelines for Keeping Experimental Animals released by the Ministry of Health of China.

Extract of *Cochinchina momordica* seeds

ECMS was prepared following the method described by Xiao *et al.* (18). The powder of *Cochinchina momordica* seeds was submerged in 50% ethanol for 24 hr. The mixture was refluxed in a round bottom flask three times at 90°C, with 2 hr for each reflux, and the ethanol was then evaporated with a R502B rotary evaporator (Shenke Tech Co. Ltd, Shanghai, China). After that, the extract was washed with diethyl ether to eliminate the substance soluble in ether. The saponin fraction was dissolved in water saturated *n*-butanol and the butanol-soluble portion was passed through a chromatography column with macroporous resin D101A (Hai Guang Chemical Co. Ltd, Tianjin, China) to remove impurities. Refined ECMS was harvested by removing the liquid eluted from the column.

FMDV vaccine and dexamethasone

FMDV vaccine was a commercial oil-adjuvanted bivalent vaccine (serotypes O and Asia I) made by Xinjiang Tiankang Animal Science Bio-Technology Co., Yinin, China. Dexamethasone sodium phosphate (DEX) (5 mg/ml) was a product of Sishui Xierkang Pharmaceutical Co., Shishui, China.

Administration of DEX, ECMS and FMDV vaccine

Fifty-six ICR mice were randomly allocated into seven groups, and each group consisted of eight animals. The mice were treated according to the schedule described in Table 1. Briefly, mice in groups 5 to 7 were s.c. injected with 0.5 mg dexamethasone daily for 4 days to induce immunosuppression. The animals were then orally given

Table 1. Treatment schedule of DEX, ECMS and FMDV vaccine to ICR mice ($n = 8/\text{group}$)

Group no.	Group	Treatment Day of administration	DEX Day 1–4	ECMS Days 5 and 26	FMDV vaccine Days 6 and 27
1	Saline		^a	— ^b	—
2	Control		/	—	+
3	ECMS (oral)		/	+ (oral)	+
4	ECMS (s.c.)		/	+ (s.c.)	+
5	DEX		+	—	+
6	DEX + ECMS (oral)		+	+ (oral)	+
7	DEX + ECMS (s.c.)		+	+ (s.c.)	+

^aMice received no treatment.

^bMice were given saline only.

ECMS (200 µg in 250 µl saline) in groups 3 and 6 or 250 µl saline in group 2, or s.c. injected with ECMS (50 µg in 100 µl saline) in groups 4 and 7 or 100 µl saline in group 5. After that, except the animals in group 1, all other animals were s.c. immunized with 100 µl oil-adjuvanted bivalent FMDV vaccine (serotypes O and Asia I). Mice in group 1 received injection of 100 µl saline only. Three weeks after the first immunization, a boosting immunization, and administration of DEX and ECMS before boosting were carried out in the same manner as described above.

Sample collection and animal observation

Blood samples were collected 2 weeks after boosting immunization for the detection of FMDV-specific IgG responses and IgG isotypes. The bodyweight of each mouse was measured on days 0, 7, 11, 32 and 46 to evaluate the effects of DEX and ECMS on the mean bodyweight of mice.

Measurement of FMDV (serotypes O and Asia I)-specific IgG and the IgG isotypes

An indirect double antibody sandwich ELISA was used for the determination of serum IgG and the isotypes as previously reported by Xiao *et al.* (18). In brief, the wells of polyvinyl 96-well microtiter plates were added to 50 µl rabbit anti-FMDV (type O or Asia I) antibody (Lanzhou Veterinary Research Institute, Lanzhou, China) in 0.05 M carbonate/bicarbonate buffer, pH 9.6 (1:1000) and incubated overnight at 4°C. After washing the wells with PBS containing 0.05% Tween-20 (PBST), they were incubated with 3% skimmed milk for blockage at 37°C for 2 hr. In the later steps, PBST was used as a diluent as well as a washing solution. Afterward, the wells were added to 50 µl FMDV antigen (LVRI) (for detection of serotype O-specific IgG, antigen was diluted 1:3 with 3% skimmed milk/PBS and for measurement of serotype Asia I-specific antibody responses, antigen was diluted 1:10 with 3%

skimmed milk/PBS) and incubated at 4°C for 2 hr. Following washing, the wells were added in duplicate with 50 µl of serum samples (1:50) and kept warm at 37°C for 1 hr. After another washing, all wells were added to 50 µl goat antimouse IgG (1:1000) (Kirkegaard, Perry Lab., Gaithersburg, MD, USA) and incubated at 37°C for 1 hr. For the determination of subclasses, 100 µl biotin-conjugated goat antimouse IgG1 or IgG2a or IgG2b or IgG3 (1:600 dilution, Santa Cruz Biotechnology Inc., Santa Cruz, CA) was added to the corresponding plate and then incubated for 1 hr at 37°C. Subsequent to washing, each well was added to 100 µl horseradish-peroxidase-conjugated anti-biotin (BD Biosciences Pharmingen, Franklin Lakes, NJ, USA) (1:4000 in PBST) and incubated for 1 hr at 37°C. One more wash was carried out and each well was added to 50 µl TMB solution (100 µg/ml 0.1 M citrate-phosphate, pH 5.0). Following development for 15 min at 37°C, 50 µl of 2 M H₂SO₄ was added to each well to stop the reaction. An automatic ELISA plate reader was used to read the optical density of the plate at 450 nm.

Statistical analysis of data

All data were subjected to one-way analysis of variance. Bonferroni method of multiple comparisons was carried out to compare the parameters among groups (22). For all the tests, $P < 0.05$ was considered significant. Data were expressed as mean \pm SD.

RESULTS

Effects of ECMS on the mean bodyweight of mice

The effects of ECMS on the bodyweight of mice are shown in Table 2. The average bodyweights of mice in dexamethasone-treated groups were significantly lower than untreated groups on day 11. Otherwise, no statistical difference was observed among groups.

Table 2. Effects of DEX and ECMS on the mean bodyweight (g; mean \pm SD) of mice ($n = 8$ /group)

Group no.	Group	Day 0	Day 7	Day 11	Day 32	Day 46
1	Saline	20.53 \pm 0.52	25.05 \pm 1.41	24.49 \pm 2.41 ^a	32.01 \pm 2.31	34.14 \pm 3.27
2	Control	20.45 \pm 1.42	23.96 \pm 1.04	25.63 \pm 1.43 ^a	30.79 \pm 2.35	32.23 \pm 3.34
3	ECMS (oral)	20.45 \pm 1.14	22.76 \pm 1.28	25.34 \pm 1.87 ^a	31.31 \pm 2.80	33.28 \pm 2.75
4	ECMS (s.c.)	20.46 \pm 0.73	25.53 \pm 1.42	27.49 \pm 1.38 ^a	31.93 \pm 1.80	31.99 \pm 4.08
5	DEX	20.35 \pm 0.72	24.26 \pm 0.81	22.79 \pm 1.07 ^b	31.34 \pm 2.35	33.09 \pm 2.26
6	DEX + ECMS (oral)	20.46 \pm 1.09	25.20 \pm 1.15	23.93 \pm 1.11 ^b	31.54 \pm 2.00	33.48 \pm 2.20
7	DEX + ECMS (s.c.)	20.35 \pm 0.92	24.09 \pm 1.86	22.85 \pm 1.81 ^b	29.88 \pm 2.80	32.38 \pm 2.95

Mean bodyweights with different superscript letters are significantly different.




Fig. 1. Serum IgG responses to FMDV serotypes O (A) and Asia I (B). Mice ($n = 8/\text{group}$) were treated with (1) saline solution; (2) FMDV antigen; (3) oral ECMS + FMDV; (4) injection ECMS + FMDV; (5) DEX + FMDV; (6) DEX + oral ECMS + FMDV and (7) DEX + injection ECMS + FMDV. After booster immunization, blood samples were collected for the measurement of FMDV-specific IgG responses by an indirect ELISA. Values are presented as mean \pm SD. Bars with different letters denote statistically significant differences ($P < 0.05$).

FMDV (serotypes O and Asia I)-specific IgG responses

The effects of ECMS on the humoral immune responses against FMDV serotype O and Asia I are depicted in Figure 1. Both oral administration of ECMS (200 μg) and s.c. injection of ECMS (50 μg) significantly ($P < 0.05$) increased serum IgG responses to FMDV serotypes O and Asia I when compared with the control (group 2 in Fig. 1). Injection of DEX suppressed IgG responses to FMDV serotype O significantly or to FMDV serotype Asia I numerically when compared with the control (group 2 in Fig. 1) but oral administration or injection of ECMS (groups 6 and 7) numerically increased IgG levels when compared with the control (group 5 in Fig. 1).

FMDV (serotypes O and Asia I)-specific IgG isotypes

The effects of ECMS on IgG isotypes IgG1, IgG2a, IgG2b and IgG3 to FMDV serotypes O and Asia I are shown in Figure 2. Oral administration and injection of ECMS tended to enhance all IgG isotype responses to immunization of bivalent FMDV vaccine (serotypes O and Asia I)

when compared with the control in which only FMDV vaccine was injected. Injection of DEX suppressed IgG isotype responses but oral administration or injection of ECMS tended to elevate IgG isotype levels.

DISCUSSION

Enhanced humoral immune response of mice to vaccination of bivalent FMDV serotypes O and Asia I vaccine has been demonstrated by oral administration or s.c. injection of an extract made from ECMS. After oral administration of ECMS (200 μg) or s.c. injection of ECMS (50 μg), immunization of a commercial FMDV vaccine induced significantly higher serum specific IgG and most IgG isotype responses than in mice given saline solution alone. In addition, giving ECMS tended to enhance serum-specific IgG and IgG isotype responses of mice immunosuppressed by s.c. injection of dexamethasone.

The mouse model has been used to study the immunity of a host against FMDV infections (23–25). Salguero *et al.* (23) have reported that mice immunized with conventional inactivated FMDV vaccine can be protected against challenge with a lethal dose of FMDV. Wong *et al.* (25) have




Fig. 2. Serum IgG isotype responses to FMDV serotypes O (A) and Asia I (B). Mice ($n = 8/\text{group}$) were treated with (1) saline solution; (2) FMDV antigen; (3) oral ECMS + FMDV; (4) Injection ECMS + FMDV; (5) DEX + FMDV; (6) DEX + oral ECMS + FMDV and (7) DEX + injection ECMS + FMDV. After booster immunization, blood samples were col-

lected for the measurement of FMDV-specific IgG isotypes IgG1, IgG2a, IgG2b and IgG3 responses by an indirect ELISA. Values are presented as mean \pm SD. Bars with different letters denote statistically significant differences ($P < 0.05$).

observed that a FMD DNA vaccine inducing an immune response in mice can also elicit protection in swine against FMD infection. Humoral immune response has been reported as an important defense mechanism against FMD virus (26, 27), and the contribution of antibodies to the major immune defense against the virus is clear (28, 29).

Studies with animal models (28, 30) have suggested that a specific humoral response is vital in the immunity against FMD virus. However, poor antibody response to FMD vaccination has been previously reported in both experimental animals and pigs (24, 31). Figure 1 shows that oral administration or s.c. injection of ECMS significantly

enhanced serum FMDV serotypes O and Asia I-specific IgG levels to a commercial bivalent FMDV vaccine. This result is consistent with our previous studies where immune responses were enhanced by co-administration of ECMS with commercial FMDV vaccine in pigs (18) or avian influenza (H5N1) vaccine in chickens (19).

Immunosuppression constitutes one of the reasons for poor immune responses to vaccination. In the present study, daily s.c. injection of DEX for 4 days effectively inhibited the immunity of mice, resulting in suppressed IgG and IgG isotype responses to FMDV vaccination (groups 5–7 on day 11 in Fig. 2). Nevertheless, oral administration or s.c. injection of ECMS caused numerically higher serum-specific IgG and IgG isotypes than in mice without ECMS treatment, indicating ECMS might have an immunomodulatory effect on immunosuppressed mice.

A conventional approach for the improvement of the efficacy of vaccination is to add adjuvant to vaccines. Adjuvant used for this purpose should be safe enough to induce minimal adverse effects to prove acceptable for use in healthy individuals. Many natural products have been reported having immunomodulatory properties, whereas their modes of action are usually unclear. Purification of the herbal extracts is usually difficult, and irritation will take place when unpurified herbal extracts are co-injected with immunizing antigens. As traditional medicinal herbs are generally given by oral route, oral use of immunomodulators can avoid the side-effects found in parenteral administration. For example, oral administration of crude saponins made from the bark of the *Quillaja* tree has been proven to have immunopotentiating activity with moderated toxicity, although they are toxic when given parenterally (32, 33). In the present study, neither abnormal behavior nor adverse side-effects were found in mice throughout the experiment, and there was no significant difference in the bodyweight between the mice given ECMS and the control mice given saline solution as indicated in Table 2 (groups 3 and 4 to group 2, groups 6 and 7 to group 5, respectively), suggesting that oral administration of ECMS is safe.

Immunosuppression constitutes one of the reasons for poor immune responses to vaccination. In this study, daily s.c. injection of DEX for 4 days effectively inhibited the immunity of mice, resulting in suppressed IgG and IgG isotype responses to FMDV vaccination (groups 5–7 on day 11 in Fig. 2). However, oral administration or s.c. injection of ECMS caused numerically higher serum-specific IgG and IgG isotypes than in mice without ECMS treatment, indicating that ECMS might have an immunomodulatory effect on immunosuppressed mice.

In summary, oral administration or s.c. injection of ECMS augmented responses of serum-specific IgG and most IgG isotypes to immunization of a commercial

FMDV (serotypes O and Asia 1) vaccine in mice. Giving ECMS tended to enhance serum IgG and IgG isotypes of mice immunosuppressed by s.c. injection of DEX. Considering the safety and immunomodulatory effect of ECMS in both normal and immunosuppressed mice demonstrated in the present study, this extract deserves further investigation to elucidate its potential in improving FMD vaccination in farm animals such as pigs, sheep and cattle.

ACKNOWLEDGMENTS

This study was supported by the National Scientific Foundation of China and the Ministry of Science and Technology of China (2008BADB4B06-2). This work is part of a doctoral program and Kedsirin Sakwiwatkul (first author) wishes to acknowledge the doctoral funds provided by Mahasarakham University, Thailand. The authors would like to thank Li-juan Zhai and Yu-tao Li for their assistance during the study.

REFERENCES

1. Belsham G.J. (1993) Distinctive features of foot-and-mouth disease virus, a member of the picornavirus family; aspects of virus protein synthesis, protein processing and structure. *Prog Biophys Mol Biol* **60**(3): 241–60.
2. Domingo E., Escarmis C., Baranowski E., Ruiz-Jarabo C.M., Carrillo E., Nunez J.L., Sobrino F. (2003) Evolution of foot-and-mouth disease virus. *Virus Res* **91**(1): 47–63.
3. Knowles N.J., Samuel A.R. (2003) Molecular epidemiology of foot-and-mouth disease virus. *Virus Res* **91**(1): 65–80.
4. Office-International-des-Epizooties. 2008. Office-International-des-Epizooties. <http://www.oie.int>.
5. He Q.G., Chen H.C., Wu B., Suo X.F., Fang L.R., Wang C. (2007) Detection of serum antibody against hog cholera, porcine parvovirus and the foot-and-mouth disease on 44 pig farms of six provinces. *Chin J Prev Vet Med* **22**(1): 5–9.
6. Wang Y.M., Lu H.L., Ying Q.X. (2007) Analysis of serum antibodies elicited by foot-and mouth disease vaccine in pigs. *Fujian Anim Husb Vet Med* **6**: 32–32.
7. Wei D.S. (2007) Discussion on the failure of vaccination in animals. *J Anim Sci Vet Med* **26**: 33–4.
8. Hao D.L., Luo R., Xu Y.F., Li K. (2005) Analysis of antibody titers to foot-and-mouth disease in pigs. *J Anim Sci Vet Med* **24**: 2–2.
9. Disis M.L., Bernhard H., Shiota F.M., Hand S.L., Gralow J.R., Huseby E.S., Gillis S., Cheever M.A. (1996) Granulocyte-macrophage colony-stimulating factor: an effective adjuvant for protein and peptide-based vaccines. *Blood* **88**(1): 202–10.
10. Jungers P., Devillier P., Salomon H., Cerisier J.E., Courouce A.M. (1994) Randomised placebo-controlled trial of recombinant interleukin-2 in chronic uraemic patients who are non-responders to hepatitis B vaccine. *Lancet* **344**(8926): 856–7.
11. Mitwalli A. (1996) Responsiveness to hepatitis B vaccine in immunocompromised patients by doubling the dose scheduling. *Nephron* **73**(3): 417–20.
12. Quiroga J.A., Castillo I., Porres J.C., Casado S., Saez F., Gracia Martinez M., Gomez M., Inglada L., Sanchez-Sicilia L., Mora A.

- (1990) Recombinant gamma-interferon as adjuvant to hepatitis B vaccine in hemodialysis patients. *Hepatology* **12**(4 Pt 1): 661–3.
13. Rey D., Krantz V., Partisani M., Schmitt M.P., Meyer P., Libbrecht E., Wendling M.J., Vetter D., Nicolle M., Kempf-Durepaire G., Lang J.M. (2000) Increasing the number of hepatitis B vaccine injections augments anti-HBs response rate in HIV-infected patients. Effects on HIV-1 viral load. *Vaccine* **18**(13): 1161–5.
14. Taglietti M. (1995) Related vaccine adjuvancy: a new potential area of development for GM-CSF. *Adv Exp Med Biol* **378**: 565–9.
15. Xu G.J. (1996) *Mu Bie Zi (Semen momordicae)*. In: Xu Q.F., ed. *Coloured Illustrations of Chinese Traditional and Herbal Drugs*, 2nd edn. Fuzhou: Fujian Science & Technology Press. p. 496.
16. Gao X.M. (2005) *Mu Bie Zi (Semen momordicae)*. In: Gao X.M., ed. *Chinese Materia Medica*. Beijing: Traditional Chinese Materia Medica Press, pp. 601–2.
17. Iwamoto M., Okabe H., Yamauchi T. (1985) Studies on the constituents of *Momordica cochinchinensis* SPRENG. II: Isolation and characterization of the root saponins, momordins I, II and III. *Chem Pharm Bull* **33**(1): 1–7.
18. Xiao C.W., Rajput Z.I., Liu D.W., Hu S.H. (2007) Enhancement of serological immune responses to foot-and-mouth disease vaccine by a supplement made of extract of cochinchina momordica seeds. *Clinical and Vaccine Immunology: CVI* **14**(12): 1634–9.
19. Xiao C.W., Hu S.H., Rajput Z.I. (2007) Adjuvant effect of an extract from *Cochinchina momordica* seeds on the immune responses to ovalbumin in mice. *Front Agric China* **1**(1): 90–5.
20. Rajput Z.I., Xiao C.W., Hu S.H., Arijo A.G., Soomro N.M. (2007) Improvement of the efficacy of influenza vaccination (H5N1) in chicken by using extract of *Cochinchina momordica* seed (ECMS). *J Zhejiang Univ Sci B* **8**(5): 331–7.
21. Baus E., Andris F., Dubois P.M., Urbain J., Leo O. (1996) Dexamethasone inhibits the early steps of antigen receptor signaling in activated T lymphocytes. *J Immunol* **156**(12): 4555–61.
22. Altman D.G. (1999) Comparing groups-continuous data. In: Altman D.G., ed. *Practical Statistics for Medical Research*. London: Chapman & Hall/CRC; pp. 179–223.
23. Salguero F.J., Sanchez-Martin M.A., Diaz-San Segundo F., de Avila A., Sevilla N. (2005) Foot-and-mouth disease virus (FMDV) causes an acute disease that can be lethal for adult laboratory mice. *Virology* **332**(1): 384–96.
24. Shi X.J., Wang B., Wang M. (2007) Immune enhancing effects of recombinant bovine IL-18 on foot-and-mouth disease vaccination in mice model. *Vaccine* **25**(7): 1257–64.
25. Wong H.T., Cheng S.C.S., Chan E.W.C., Sheng Z.T., Yan W.Y., Zheng Z.X. (2000) Plasmids encoding foot-and-mouth disease virus VP1 epitopes elicited immune responses in mice and swine and protected swine against viral infection. *Virology* **278**(1): 27–35.
26. Brown F. (1995) Antibody recognition and neutralization of foot-and-mouth disease. *Semin Virol* **6**: 243–8.
27. Meloen R.H., Rowlands D.J., Brown F. (1979) Comparison of the antibodies elicited by the individual structural polypeptides of foot-and-mouth disease and polio viruses. *J Gen Virol* **45**: 761–3.
28. McCullough K.C., Crowther J.R., Butcher R.N., Carpenter W.C., Brocchi E., Capucci L., De Simone F. (1986) Immune protection against foot-and-mouth disease virus studied using virus-neutralizing and non-neutralizing concentrations of monoclonal antibodies. *Immunol* **58**(3): 421–8.
29. McCullough K.C., De Simone F., Brocchi E., Capucci L., Crowther J.R., Kihm U. (1992) Protective immune response against foot-and-mouth disease. *J Virol* **66**(4): 1835–40.
30. McCullough K.C., Parkinson D., Crowther J.R. (1988) Opsonization enhanced phagocytosis of foot-and-mouth disease virus. *Immunol* **65**: 187–91.
31. Xiao C.W., Rajput Z.I., Hu S.H. (2007) Improvement of a commercial foot-and-mouth disease vaccine by supplement of Quil A. *Vaccine* **25**(25): 4795–4800.
32. Chavali S.R., Campbell J.B. (1987) Adjuvant effects of orally administered saponins on humoral and cellular immune responses in mice. *Immunobiology* **174**(3): 347–59.
33. Maharaj I., Froh K.J., Campbell J.B. (1986) Immune responses of mice to inactivated rabies vaccine administered orally: potentiation by Quillaja saponin. *Can J Microbiol* **32**(5): 414–20.