Contents lists available at ScienceDirect

Food and Chemical Toxicology

journal homepage: www.elsevier.com/locate/foodchemtox



Toxicity of Jatropha curcas phorbol esters in mice

Cai-Yan Li^a, Rakshit K. Devappa^b, Jian-Xin Liu^{a,*}, Jian-Min Lv^c, H.P.S. Makkar^{b,*}, K. Becker^b

^a College of Animal Science, Zhejiang University, 310029 Hangzhou, PR China

^b Institute for Animal Production in the Tropics and Subtropics, University of Hohenheim, 70593 Stuttgart, Germany

^c Laboratory Animal Research Center, Zhejiang Chinese Medicine University, 310053 Hangzhou, PR China

ARTICLE INFO

Article history: Received 16 July 2009 Accepted 19 November 2009

Keywords: Phorbol esters Toxicity Jatropha curcas LD₅₀ Mice Histopathology

ABSTRACT

Phorbol esters are the main toxins in *Jatropha curcas* seed and oil. The aim of this study was to assess the acute toxicity of phorbol esters given by intragastric administration and to determine the LD_{50} for Swiss Hauschka mice. The LD_{50} and 95% confidence limits for male mice were 27.34 mg/kg body mass and 24.90–29.89 mg/kg body mass; and the LD_5 and LD_{95} were 18.87 and 39.62 mg/kg body mass, respectively. The regression equations between the probits of mortalities (*Y*) and the log of doses (*D*) was $Y = -9.67 + 10.21 \log (D)$. Histopathological studies on the organs from the dead mice showed: (1) no significant abnormal changes in the organs at the lowest dose (21.26 mg/kg body mass) studied, (2) prominent lesions mainly found in lung and kidney, with diffused haemorrhages in lung, and glomerular sclerosis and atrophy in kidney at doses ≥ 32.40 mg/kg body mass, and (3) multiple abruption of cardiac muscle fibres and anachromasis of cortical neurons at the highest dose of 36.00 mg/kg body mass. The results obtained would aid in developing safety measures for the Jatropha based biofuel industry and in exploiting the pharmaceutical and agricultural applications of phorbol esters.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Jatropha curcas is a multipurpose bush or a small tree belonging to the family of *Euphorbiaceae*. It is a native of tropical America, but now thrives in many parts of the tropics and sub-tropics in Africa, Asia and southern America (Gübitz et al., 1999). The plant is well adapted to barren or drought affected areas and even survives in poor stony soils. *J. curcas* seed contains high amount of oil that can be converted into biodiesel of high quality upon transesterification. Apart from the oil, the seed cake or kernel meal leftover has gained tremendous interest for their utilization in feed formulations (Makkar et al., 1997). *J. curcas* kernel meal is rich in protein and the essential amino acid composition of the protein, except lysine, is comparable to that of soybean meal (Makkar et al., 1998; Devappa and Bhagya, 2008). However, the main toxins present in these by-products are phorbol esters, which prevent their utilization as feed ingredients (Makkar et al., 1997).

Phorbol esters are diterpenes having tigliane skeletal structure. Six phorbol esters have been characterized from *J. curcas* (Haas et al., 2002). Phorbol esters are amphiphilic molecules and have tendency to bind phospholipid membrane receptors. During normal signal transduction process, DAG (diacyl glycerol) activates PKC (protein kinase C), which is involved in various other signal transduction pathways. The phorbol esters act as an analogue for DAG and are strong PKC activators. These hyperactivate PKC and trigger cell proliferation, thus amplifying the efficacy of carcinogens. Phorbol esters themselves do not induce tumors but promote tumor growth following exposure to subcarcinogenic dose of carcinogen. They can thus be regarded as co-carcinogens (Goel et al., 2007). The concentration of phorbol esters varies from 2 to 3 mg/g in kernel and 2 to 4 mg/g in oil in different varieties of *J. curcas* (Makkar et al., 1997). Although phorbol esters are lipophilic, they get strongly bound to the matrix of kernel meal. Approximately 30% of the total phorbol esters were present in kernel meal containing <1% lipid (Makkar et al., 2008).

Studies in the last decade showed that *J. curcas* exhibited toxicity to a wide variety of organisms, from microorganisms to higher animals. But, the toxicity effects studied in animals are mainly by force-feeding either *J. curcas* raw or defatted seed meals, leaves or their various organic solvent/aqueous extracts. Oral administration of Jatropha kernel oil obtained by hexane–soxhlet extraction had an LD₅₀ of 6 ml/kg body mass in rats. The rats, at 6, 9 and 13.5 ml/kg body mass exhibited diarrhoea and haemorrhagic eyes; and the autopsy showed inflammation of the gastro-intestinal tract (Gandhi et al., 1995). *J. curcas* oil at a dose of 2 g/kg body mass caused significant acute toxicity by inhibiting the birth of pups in rats (Odusote et al., 2002). The concentrated toxic fraction obtained after successive extraction of oil by 90% methanol and ethyl ether, as reported by Gandhi et al. (1995) exhibited toxicity



^{*} Corresponding authors. Tel.: +49 71145923640; fax: +49 71145923702 (H.P.S. Makkar).

E-mail addresses: liujx@zju.edu.cn (J.-X. Liu), makkar@uni-hohenheim.de (H.P.S. Makkar).

towards rabbit (100 µl), mice and rats (50 µl). The common symptoms of topical application were erythema, oedema, necrosis, diarrhoea, scaling and thickening of the skins (Gandhi et al., 1995). Feeding of J. curcas seeds, fruits or leaves caused toxicity depending on the dose and the animal model tested. Raw or defatted seeds when force-fed to fish, chicks, pigs, goat, mice and rats (caused severe toxicity symptoms before death (Liberalino et al., 1988; Chivandi et al., 2000; Gadir et al., 2003; Adam and Magzoub, 1975; Adam, 1974). Various organic and aqueous extracts also exhibited different toxic symptoms depending on dose, mode of administration and sensitivity of the animals being tested (Trebien et al., 1988; Mariz et al., 2006, 2008). For example, acetonitrile extract of J. curcas (seed or oil) when given to albino rats at an oral dose of 50 mg/kg body mass (single dose) produced mild toxicological, biochemical and histopathological changes (Abd-Elhamid, 2004). The methanol, petroleum ether and dichloromethane extracts of I. curcas fruit caused foetal resorption indicating pregnancy terminating effect in rats (Goonesekera et al., 1995). The irritant methanol fraction from J. curcas oil induced tumor promotion upon topical initiation by 7, 12-dimethylbenz(a)anthracene (DMBA) in mice, with 36% of the animals having skin tumors in 30 weeks (Horiuchi et al., 1988).

Hitherto, all the LD_{50} studies, toxicity studies or the histopathological studies conducted were using either *J. curcas* oil, seed, defatted seed cake or crude fractions from them. In the present study we purified phorbol esters from *J. curcas* oil, evaluated their acute toxicity when administered intragastrically and determined LD_{50} for laboratory mice.

2. Materials and methods

2.1. Materials

The *J. curcas* oil used in the study was from toxic genotypes from India. Corn oil was obtained from Sigma Chemical Company (Darmstadt, Germany). All other chemicals were of analytical grade.

2.2. Phorbol esters isolation and quantification

Phorbol esters were determined in quadruplicate according to Makkar et al. (2007), which was based on the method of Makkar et al. (1997). Briefly, 0.5 g of the sample was extracted four times with methanol. A suitable aliquot was loaded on a high-performance liquid chromatography (HPLC) reverse-phase C18 LiChrospher 100, 5 μ m (250 \times 4 mm I.D. from Merck (Darmstadt, Germany) column. The column was protected with a head column containing the same material. The separation was performed at room temperature (23 °C) and the flow rate was 1.3 ml/min using a gradient elution (Makkar et al., 2007). The four-phorbol esters peaks were detected at 280 nm and appeared between 25.5 and 30.5 min. For LD_{50} studies, the phorbol esters were carefully collected at the above retention times. All the collected fractions were in approximately 90% acetonitrile. These fractions were pooled and kept in a freezer (-20 $^\circ$ C). To avoid oxidation by water, the top acetonitrile layer from the frozen sample was separated and the acetonitrile was rotary evaporated at low vacuum to collect the colorless oily fraction. This fraction was redissolved in methanol and subjected to HPLC for checking the purity and concentration. The results were expressed as equivalent to phorbol-12-myristate 13-acetate.

Table 1	
Mortality of mice after	intragastric administration.

2.3. Test substance

Phorbol esters were isolated from the *J. curcas* oil as above and diluted in high purity corn oil (Acros Organics Co., Geel, Belgium) for oral administration.

2.4. Animals and conditions

All the conditions used for animal housing and handling were approved by Animal Care and Use Committee at Zhejiang Province (China), and the experimental protocols used followed the Regulations for the Administration of Affairs Concerning Experimental Animals (The State Council of People's Republic of China, 1988).

A total of 70 male specific pathogen-free (SPF) grade Swiss Hauschka mice (21 days old) were obtained from Shanghai Laboratory Animal Center of China Academy of Science (Shanghai, China). The number of laboratory-animal-quality certification was SCXK (Hu) 2003-0003. The mice with an initial body mass of 17–18 g were kept in a large clean $27 \times 17 \times 13$ cm polycarbonate cages (5 mice per cage) with bedding and fed on a laboratory diet (radiated by Co⁶⁰) for 3 days for acclimatization. After acclimatization, the mice were weighed and numbered.

The experiment was conducted in a barrier system with an experimental facility (License No: SYXK (ZHE) 2003-0003). The housing conditions were controlled automatically with a room temperature of 22 ± 1 °C; relative humidity 50–70%; lighting 150–200Lx, the sequence being 12 h dark and light cycle; noise <50 dB; air cleanliness degree 10,000 grade. During the experiment the mice had *ad libitum* access to food and sterile drinking water.

2.5. Test procedure

The test procedures were in accordance with the Regulatory Guide on the Techniques for Drug Research (The State Food and Drug Administration of People's Republic of China, 2005). A pretest was conducted to observe the range of toxicity so that the proper dose levels could be established for LD_{50} determination. Three dose levels (6, 12 and 18 mg/kg body mass) of phorbol esters were used for the pretesting. Based on the pretest results, six dosages (36.00, 32.40, 29.16, 26.24, 23.62 and 21.26 mg/kg body mass) were established with each group comprised of 10 mice using random block design (average body mass 18–20 g). The mice were kept fasting for 12 h before the phorbol esters doses were given by intragastric administration. The dosage of phorbol esters was given to mice as 0.2 ml/10 g body mass. The remaining 10 mice served as control and were given an equal volume (0.2 ml/10 g body mass) of corn oil. After the oil or phorbol esters administration, the mice in each group were fed as normal, and had free access to food and drinking water.

In the following 19 days, the clinical signs, change in body mass, and toxicity symptoms in mice were observed every day.

2.6. Histopathological studies

All deaths were recorded. Died mice and mice killed by cervical dislocation at the end of experiment were examined for gross and microscopical changes. The specimens of tissues from kidneys, liver, heart, spleen, brain and the lungs were taken for histopathological examination. The tissues were immediately rinsed with physiological saline, fixed overnight in 4% paraformaldehyde, and then dehydrated in a graded series of ethanol and embedded in paraffin for later slices and haematoxylin and eosin staining.

2.7. Calculations and statistical analysis

The LD_{50} determination was based on dose levels that increased by a geometrical progression. Six dose levels were ultimately required for establishing the LD_{50} . Calculations for LD_{50} , 95% confidence limits, LD_5 and LD_{95} were based on the Bliss method as explained in Zhou (1988). The Bliss was calculated by using the NDST Software Version 8.0 (Sun, 1998).

Dose (mg/kg)	Dead	After administration (days)														
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15-19
0 (Control)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
36.00	9	0	1	0	2	2	3	0	0	0	0	0	0	0	1	0
32.40	7	0	2	2	1	1	0	0	0	0	0	1	0	0	0	0
29.16	7	0	1	2	2	0	0	1	1	0	0	0	0	0	0	0
26.24	4	0	0	1	0	0	2	0	1	0	0	0	0	0	0	0
23.62	3	0	0	1	1	0	1	0	0	0	0	0	0	0	0	0
21.26	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0

Т	able 2	
C	hanges in total body mass (g) of mice after administration (Mean ± SD)	1.

Dose (mg/kg)	Body mass (g) after administration (days)							
	1	3	5	7	14	19		
36.00	19.0 ± 0.7	17.6 ± 1.8	17.2 ± 3.5	21.1 ± 5.9	28.4	30.60		
32.40	19.1 ± 0.8	17.0 ± 1.1	15.9 ± 2.0	17.9 ± 2.6	23.8 ± 1.7	27.6 ± 0.7		
29.16	19.8 ± 0.9	17.9 ± 1.2	16.3 ± 2.3	17.2 ± 4.0	23.8 ± 3.9	28.1 ± 2.2		
26.24	18.9 ± 0.8	17.2 ± 0.8	16.8 ± 2.2	18.2 ± 3.3	25.0 ± 2.1	28.3 ± 1.8		
23.62	19.5 ± 0.7	17.6 ± 0.7	16.1 ± 1.9	17.0 ± 2.2	22.1 ± 4.7	26.6 ± 6.4		
21.26	18.8 ± 0.7	17.1 ± 1.4	16.7 ± 2.6	18.4 ± 3.5	24.4 ± 3.0	28.3 ± 2.2		
0 (control)	19.2 ± 0.8	22.6 ± 1.1	23.9 ± 1.4	25.9 ± 1.6	27.5 ± 1.7	29.8 ± 1.9		

All the experimental analysis were statistically analyzed using the General Linear Model (GLM) Procedure of SAS by Duncan's new multiple range test to determine the significant differences in each concentration doses (p < 0.05).

3. Results

3.1. Mortality and live weight changes

The starting material had a phorbol ester concentration of 45.03 mg/g in corn oil. The death of mice occurred in a dose dependent manner (Table 1). Mice began to die on the 2nd day after administering the phorbol esters, and no more mice died during the extended 15–19 observation days. The time to death seems to depend on the administered dose and the individual susceptibility of mice to the phorbol esters. The higher the dose or less the resistance of an individual, the quicker the animal died.

Changes in body mass after administration of phorbol esters are shown in Table 2. For all doses, body mass of the phorbol esters administered mice decreased significantly after 3 days of the administration (p < 0.05). However, at the doses of 21.26–32.40 mg/kg, the body mass of mice those survived for 7 days after phorbol esters administration started reverting back to normal. At the highest dose of 36.00 mg/kg, the recovery of body weight occurred after 5 days of phorbol esters administration.

3.2. LD₅₀ determinations

Phorbol esters LD_{50} and its 95% confidence limits for male mice were 27.34 mg/kg body mass and 24.90–29.89 mg/kg body mass; and LD_5 and LD_{95} were 18.87 and 39.62 mg/kg body mass, respectively (Table 3). The regression equation between the probits of mortalities (Y) and the log of doses (D) was derived as: Y(probit) = $-9.67 + 10.21 \log (D)$.

3.3. Clinical signs and histopathological findings

The mice administrated with phorbol esters showed varied degrees of toxicity reactions at different doses. At the highest dose of 36.00 mg/kg, the affected animals showed depression, languishment, loss of body mass, closing of eyes, humidity of anus, arch of the back and behaviors of being easily affrighted and crouched together. For the mice administrated with phorbol esters at all doses, the stool in rectum was dry in the form of beads, and both small and large intestines were filled with black digesta. In contrast, the stool from the control mice was wet and soft. The forming of dry bead like structure in the rectum of affected animals may be related to the intestinal dehydration, most probably caused by phorbol ester mediated electrolyte imbalance. It is supposed that

Та	ble	3

LD₅₀ determinations of phorbol esters for mice.

Dose mg/kg	Log Dose	Tested	Dead	Tested/dead (%)	Experimental probability (Y)	Regression probability (Y)
36.00	1.5563	10	9	90	6.2817	6.2203
32.40	1.5105	10	7	70	5.5240	5.7531
29.16	1.4648	10	7	70	5.5240	5.2859
26.24	1.4190	10	4	40	4.7471	4.8187
23.62	1.3733	10	3	30	4.4760	4.3515
21.26	1.3275	10	1	10	3.7183	3.8843



Fig. 1. Pathological changes of liver observed at a dose of 36.00 mg/kg (A) Fatty vacuoles in the live cells (B) Normal liver.



Fig. 2. Pathological changes of spleen observed at a dose of 36.00 mg/kg (A) Widely hyperemia and exudates (B) Normal spleen.

the gastro-intestinal tract haemorrhage may result in the black digesta in intestines. The stool findings showed that the digestive system of mice is sensitive to the presence of phorbol esters.

Histopathological studies of the organs showed normal cellular architecture at the lowest dose (21.26 mg/kg). At a dose of 23.61 mg/kg, no evidence of damage to heart, small intestine, brain and spleen was seen; however, sporadic infiltrated lymphocyte in the liver, seldom haemorrhage in lung and a few of glomerular sclerosis appeared. At a dose of 26.24 mg/kg, congestion of sinus hepaticus, haemorrhage of spleen and a few of glomerular atrophy were seen. At a dose of 29.31 mg/kg, congestion of the pulmonary alveolar capillaries was observed. When the dose increased to 32.40 mg/kg, much more pathological symptoms were seen in lung, with more mice showing congestion of the pulmonary alveolar capillaries and a few of mice exhibiting haemorrhage and burst of alveolus. At the highest dose of 36.00 mg/kg, multiple abruption of cardiac muscle fibres and anachromasis of cortical neurons appeared. Other histopathological changes included the frequent appearance of fatty vacuoles in the liver cells (Fig. 1), widely hyperemia and exudate in spleen (Fig. 2), diffuse haemorrhage and exudate in lung (Fig. 3), and glomerular sclerosis (Fig. 4). Overall, the prominent pathological symptoms were mainly observed in lung and kidney.

4. Discussion

Among the different methods available for LD_{50} determinations, the sensitivity of the Bliss procedure used in the present study is high, despite its complexity (Li et al., 1995).



Fig. 3. Pathological changes of lung observed at a dose of 36.00 mg/kg (A) Widely diffuse haemorrhage and exudates (B) Normal lung.



Fig. 4. Pathological changes of kidney observed at a dose of 36.00 mg/kg (A) Glomerular sclerosis (B) Normal kidney.

Table 4	
Influence of phorbol ester in ruminants force-fed on powdered Jatropha curcas decorticated seeds (kerne	ls).

Animal	Dose (g/kg body mass/ day)	100% Mortality (in days)	Calculated value of mg phorbol esters consumed/kg body mass ^b	References
Nubian goat	10	2-3	30-45	Adam and Magzoub
	10 ^a	6	15	(1975)
	5	3-4	22.5-30	
	1	7–9	10.5–13.5	
	0.5	11-12	8.25–9.0	
	0.25	18–21	6.75–7.87	
Nubian goat	1	6-7	9–10.5	Ahmed and Adam (1979)
	0.5	16–22	15-20.6	
	0.05	19–25	1.4–1.9	
Desert sheep	1	3–5	4.5–7.5	
	0.5	7–10	5.2–7.5	
Nubian goat kids	1	7–11	10.5–16.5 ^c	Gadir et al. (2003)
	0.25	18-21	6.75–7.87 ^c	

^a Single dose.

^b The average concentration of phorbol esters (Egyptian genotype; 1.5 mg/g, *n* = 6) analyzed in our laboratory was taken for calculation.

^c The average concentration of phorbol esters (toxic genotype; 2.5 mg/g, n = 8) analyzed in our laboratory was taken for calculation.

The LD₅₀ study indicates that the purified phorbol esters isolated from the oil are highly toxic to mice and produce severe pathological symptoms. These results support the findings of Makkar et al. (1997, 1998) that phorbol esters are the main toxins in J. curcas oil and seeds. Phorbol esters are present in leaves, stems, flowers and roots of J. curcas (Makkar and Becker, 2009) and therefore the consumption of *J. curcas* in any form, oil, seeds, seed cake or extracts is toxic to animals, and elicits severe pathological symptoms. In the ruminants, force-feeding (drenching) studies using decorticated J. curcas seeds (kernels) caused acute toxicity with 100% mortality depending on the dose administered (Table 4). This highlights the importance of complete removal of phorbol esters from the Jatropha meal before using it in feed formulations. In many parts of the world, J. curcas is used as a live fence and the presence of phorbol esters in different parts of the plant and their toxicity is responsible for this use of the plant.

Taking average phorbol esters concentration, determined in our laboratory in the seeds of a number of toxic Indian variety of *J. curcas*, we back calculated the LD_{50} reported for oil by Gandhi et al. (1995) and for methanol extract from oil reported by Oluwole and Bolarinwa (1997). The calculated oral LD_{50} is 19.19 mg/kg body mass (analyzed average value of 3.5 mg/g taken for calculation) in rats and 49.87 mg/kg (intraperitoneal) in mice, respectively (analyzed average value of 1.98 mg/g taken for calculation). Rats fed with diet containing defatted whole seed Jatropha meal (10% protein replacement level) caused severe pathological symptoms and death occurred at a phorbol esters concentration of 47.31 mg/kg body mass (Rakshit et al., 2008).

The symptoms and toxicity of *J. curcas* depend on extract, dose, mode of administration and sensitivity of the animals being tested. For example, topical application of petroleum ether extract of *J. curcas* oil (at a dose of 100 μ l) on shaved dorsal skin of rabbit showed erythema and oedema, which later became necrotic and regenerated. In mice, the same extract upon topical application (at a dose of 50 μ l) exhibited swelling of the face, haemorrhagic eyes, diarrhoea and skin erythema before death. Whereas, rats (at a dose of 50 μ l) showed oedema and erythema at 4 h of topical application on the shaved dorsal skin which subsequently led to severe scaling and thickening of the skin (Gandhi et al., 1995).

All the feeding studies on *J. curcas* showed severe clinical and pathological symptoms. Among the important symptoms observed was transient loss of body mass and mild to severe macroscopic/microscopic changes in the kidney, lungs, heart, liver, and spleen in a dose dependent manner (Adam, 1974; Liberalino et al., 1988; Rakshit et al., 2008). In the present study as well, the major pathological symptoms were observed in lung and kidney, with

widely diffuse haemorrhage in lung and glomerular sclerosis in the kidney.

The data obtained in this study would aid in developing: (i) emergency procedures in case of major spills or accidental contacts, (ii) guidelines for the use of appropriate safety clothing and equipment, and (iii) transport regulations. It would also help in establishing occupational exposure limits and in developing material safety data sheets for various by-products obtained during biodiesel production from *J. curcas*.

Conflict of Interest

The authors declare that there are no conflicts of interest.

Acknowledgements

The authors are grateful to the Bundesministerium für Bildung und Forschung (BMBF), Berlin, Germany and Ministry of Science and Technology (MoST), Beijing, China for the financial supporting.

References

- Abd-Elhamid, H.F., 2004. Investigation of induced biochemical and histopathological parameters of acetonitrile extract of *Jatropha curcas* in albino rats. J. Egypt. Soc. Parasitol. 34 (2), 397–406.
- Adam, S.E.I., 1974. Toxic effects of Jatropha curcas in mice. Toxicology 2, 67-76.
- Adam, S.E.I., Magzoub, M., 1975. Toxicity of Jatropha curcas for goats. Toxicology 4 (3), 347–354.
- Ahmed, O.M., Adam, S.E., 1979. Toxicity of Jatropha curcas in sheep and goats. Res. Vet. Sei. 27 (1), 89–96.
- Chivandi, E., Makuza, S.M., Erlanger, K.H., Mtimuni, J.P., Read, J.S., Tivapasi, M., 2000. Effects of dietary *Jatropha curcas* on the haematology of weaned pigs. Zimbabwe Vet. J. 31 (4), 83–91.
- Devappa, R.K., Bhagya, S., 2008. Biochemical and nutritional evaluation of Jatropha protein isolate prepared by steam injection heating for reduction of toxic and antinutritional factors. J. Sci. Food Agric. 88, 911–919.
- Gadir, A., Onsa, T.O., Ali, W.E.M., El Badwi, S.M.A., Adam, S.E.I., 2003. Comparative toxicity of Croton macrostachys, Jatropha curcas, Piper abyssinica seeds in Nubian goats. Small Ruminant Res. 48 (1), 61–67.
- Gandhi, V.M., Cherian, K.M., Mulky, M.J., 1995. Toxicological studies on Ratanjyot oil. Food Chem. Toxicol. 33 (1), 39–42.
- Goel, G., Makkar, H.P.S., Francis, G., Becker, K., 2007. Phorbol esters: structure, biological activity and toxicity in animals. Int. J. Toxicol. 26, 279–288.
- Goonesekera, M.M., Gunawardana, V.K., Jayasena, K., Mohammed, S.G., Balasubramaniam, S., 1995. Pregnancy terminating effect of *Jatropha Curcas* in rats. J. Ethnopharmacol. 47, 117–123.
- Gübitz, G.M., Mittelbach, M., Trabi, M., 1999. Exploitation of the tropical oil seed plant Jatropha curcas L. Bioresource Technol. 67, 73–82.

Haas, W., Strerk, H., Mittelbach, M., 2002. Novel 12 deoxy-16-hydroxyphorbol diesters isolates from the seed oil of *Jatropha curcas*. J. Nat. Prod. 65, 1434–1440.

Horiuchi, T., Suttajit, M., Suguri, H., Endo, Y., Shudo, K., Wongchai, V., Hecker, E., Fujiki, H., 1988. A new tumor promoter from the seed oil of *Jatropha curcas* L., an intramolecular diesters of 12-deoxy-16-hydroxyphorbol. Cancer Res. 48 (20), 5800–5804.

Li, Q.X., Wang, H., Xiao, Q.Q., Kong, R., 1995. The evaluation and calculation of Median Lethal Dose $(\rm LD_{50})$ using Bliss method. J. Math. Med. 4, 318–320.

- Liberalino, A.A.A., Bambirra, E.A., Moraes-Santos, T., Viera, C.E., 1988. Jatropha curcas L. seeds. Chemical analysis and toxicity. Braz. Arch. Biol. Technol. 31, 539–550.
- Makkar, H.P.S., Aderibigbe, A.O., Becker, K., 1998. Comparative evaluation of nontoxic and toxic varieties of *Jatropha curcas* for chemical composition, digestibility, protein degradability and toxic factors. Food Chem. 62 (2), 207– 215.
- Makkar, H.P.S., Becker, K., 2009. Jatropha curcas an exciting crop for generation of biofuel and value-added products. Eur. J. Lipid Sci. Technol. 11 (8), 773–787.
- Makkar, H.P.S., Becker, K., Sporer, F., Wink, M., 1997. Studies on nutritive potential and toxic constituents of different provenances of *Jatropha curcas*. J. Agric. Food Chem. 45, 3152–3157.
- Makkar, H.P.S., Martinez-Herrera, J., Becker, K., 2008. Variations in seed number per fruit, seed physical parameters and contents of oil, protein and phorbol ester in toxic and non-toxic genotypes of *Jatropha curcas*. J. Plant Sci. 3 (4), 260–265.
- Makkar, H.P.S., Siddhuraju, P., Becker, K., 2007. A Laboratory Manual on Quantification of Plant Secondary Metabolites. Human Press, Totowa, New Jersey. p. 130.
- Mariz, S.R., Arauojo, W.C., Cerqueira, G.S., Araujo, W.C., Duarte, J.C., Diniz, M.F.F.M., Medeiros, I.A., 2008. Avaliacao histopatologica em ratros apos tratamento

agudo com o extrato etanolico de partes aereas de *Jatropha gossypiifolia* L. Braz. J. Pharmacog. 18 (2), 213–216.

- Mariz, S.R., Cerqueira, G.S., Arauojo, W.C., Duarte, J.C., Melo, A.F.M., Santos, H.B., Oliveria, K., Diniz, M.F.F.M., Medeiros, I.A., 2006. Estudo toxicologixo agudo do extrato ethanolico de partes aereas de *Jatopha gossypiifolia* L. em ratos. Braz. J. Pharmacog. 16 (3), 372–378.
- Odusote, O.M., Abioye, A.O., Rotib, M.O., 2002. *Jatropha curcas* seed oil Linn (Euphorbiaceae) contraceptive activity and on oral formulation. Nig. Quart. J. Hosp. Med. 12 (1–4), 44–47.
- Oluwole, F.S., Bolarinwa, F., 1997. *Jatropha curcas* extract causes anemia in rat. Phytother. Res. 11, 538–539.
- Rakshit, K.D., Darukeshwara, J., Rathina Raj, K., Narasimhamurthy, K., Saibaba, P., Bhagya, S., 2008. Toxicity studies of detoxified Jatropha meal (*Jatropha curcas*) in rats. Food Chem. Toxicol. 46, 3621–3625.
- Sun, R.Y., 1998. NDST (New Drug Statistic) Version 8.0. Wan-Nan Medical College, Wuhu, China.
- The State Food and Drug Administration of People's Republic of China, 2005. Regulatory Guide on the Techniques for Drug Researches. Chinese Medical Science and Technology Press, Beijing. pp. 83–93.
- Trebien, H.A., Neves, P.C.A., Yunes, R.A., Calixto, J.B., 1988. Evaluation of pharmacological activity of crude hydroalcoholic extract from *Jatropha elliptica*. Phytother. Res. 2 (3), 115–118.
- Zhou, H.J., 1988. Statistical Methods for Biological Test. People's Medical Publishing House, Beijing. p. 214.