

Acta Biochim Biophys Sin, 2015, 47(2), 80–90 doi: 10.1093/abbs/gmu119 Advance Access Publication Date: 22 December 2014 Original Article

OXFORD

Original Article

Proteomic analysis of silkworm midgut cellular proteins interacting with the 5' end of infectious flacherie virus genomic RNA

Mingqian Li, Xinyi He, Han Liu, Zhangwuke Fu, Xiangkang He, and Xingmeng Lu*

Silkworm Pathology and Disease Control Laboratory, College of Animal Sciences, Zhejiang University, Hangzhou 310058, China

*Correspondence address. Tel/Fax: +86-571-88982305; E-mail: xmlu@zju.edu.cn

Received 1 August 2014; Accepted 21 October 2014

Abstract

The flacherie disease in the silkworm is caused by the infectious flacherie virus (IFV). IFV relies on its 5' region of genomic RNA to recruit host-related factors to implement viral translation and replication. To identify host proteins bound to the 5'-region of IFV RNA and identify proteins important for its function, mass spectrometry was used to identify proteins from silkworm midgut extracts that were obtained using RNA aptamer-labeled 5' region of IFV RNA. We found 325 protein groups (unique peptide ≥ 2) bound to the 5' region of IFV RNA including translation-related factors (16 ribosomal subunits, 3 eukaryotic initiation factor subunits, 1 elongation factor subunit and 6 potential internal ribosome entry site *trans*-acting factors), cytoskeleton-related proteins, membrane-related proteins, metabolism enzymes, and other proteins. These results can be used to study the translation and replication related factors of IFV interacting with host silkworm and to control flacherie disease in silkworm.

Key words: infectious flacherie virus, RNA-binding protein, mass spectrometry, silkworm, midgut cellular protein

Introduction

Bombyx mori infectious flacherie virus (IFV) is the causative agent of flacherie disease of the silkworm, which is a viral disease that causes serious sericulture losses in China, Japan, and other regions [1]. IFV is a positive-sense single-stranded RNA virus and a species of the family Iflaviridae, genus *Iflavirus* that encodes a single polyprotein of 3085 amino acids that is proteolytically processed into four structural proteins and a series of non-structural proteins by cellular and viral proteases [2]. The genomes of small positive-strand RNA viruses not only encode the viral proteins required for viral genome replication and infectious virus production, but also contain *cis*-acting RNA sequences and structures that play critical roles in the virus life cycle [3]. Highly conserved secondary RNA structures are known to be present at the 5' region of the genomes of picornaviruses and hepatitis C virus (HCV) [4,5].

Host cell factors that interact with these RNA elements play important roles in all aspects of the virus life cycle [6,7]. Translation initiation of all picornavirus RNAs is dependent on the internal ribosome entry site (IRES) located in the long 5' untranslated region (UTR) [8]. Various structural elements in the 5' UTR, which differ among picornavirus genera, control the viral replication cycle in concerted action with the 3' UTR [9,10]. RNA–protein complexes are formed when authentic viral RNA or *in vitro*-synthesized viral RNA transcripts are incubated with cell extracts. These complexes are involved in viral RNA replication and translation. RNA–protein interactions have been determined using different approaches, such as ultraviolet cross-linking, and electrophoretic mobility-shift assays. Although these techniques are useful for analyzing RNA-binding factors, mass spectrometry (MS) offers the advantage of identifying several proteins within a complex mixture [11–13]. RNA aptamers are short nucleic acids (typically 12–80 nucleotides) capable of specific, tight binding to their target molecules. They may be used as RNA affinity tags for the rapid purification and investigation of RNAs and RNA–protein complexes [14]. At present, many viral RNA-interacting cellular proteins have been identified by riboproteomic affinity methods, such as several ribosomal subunits, eukaryotic initiation factors (eIFs), and IRES *trans*-acting factors (ITAFs) [10,15].

The target of IFV is the goblet cells of the midgut epithelium, and IFV is replicated in the cell cytoplasm. Without a susceptible cell line for IFV infection, it is difficult to study the biology of virus [16]. However, this problem prompted the development of alternative strategies to generate *in vitro* RNAs that are similar to the authentic viral genome. The RNA of IFV can also act as efficient mRNA in rabbit reticulocyte lysate and in wheat germ translation systems [17]. A previous study has revealed that the 5' region of IFV has IRES activity, like *Ectropis obliqua* virus, *Perina nuda* virus, and *Varroa destructor* virus-1 in *Iflavirus* [18–21]. Interaction with these eIFs and ITAFs is necessary to promote IRES activity. The IRES region is a superior antivirus target. However, there were no reports regarding the interactions of host proteins with the 5' region of *Iflavirus*.

In this study, we incubated the 5' region of IFV genomic RNA (599 nt) that links the streptavidin aptamers [22] and the midgut extracts to perform affinity chromatography with streptavidin beads. The affinity of purified RNA–protein complexes was analyzed by MS. Our results demonstrated that the 5' end of the IFV genome contains elements that bind specifically to different midgut cellular proteins, some of which include ribosomal protein, eIF, eukaryotic translation elongation factor (eEF), and others that are known or unknown factors involved in picornavirus and HCV IRES-associated translation and replication. These results would be useful for further study of the translation and replication and replication of IFV and control of flacherie disease in silkworm.

Materials and Methods

Construction of plasmids and *in vitro* **RNA transcription** A cDNA clone of nt 1 to 599 of the 5'-region of IFV-CHN01 genome (GenBank: EU868609) was fused to the SP6 promoter and the RNA streptavidin aptamer and linker sequence using splice overlap extension polymerase chain reaction (PCR) (Fig. 1A) [22]. The PCR product was purified by a PCR purification kit (AXGEN, New York, USA) according to the manufacturer's instructions and inserted into plasmid vector pMD18 (TaKaRa, Dalian, China) linearized by the restriction enzyme *Xba*I. The linearized vectors were used for SP6 RNA polymerase (TaKaRa) *in vitro* transcription reactions to generate RNA according to the manufacturer's instructions.

Preparation of midgut cellular extracts

The midguts of 5th instar silkworm (strain P50) were collected, washed in physiologic saline and phosphate-buffered saline (PBS), and homogenized with insect cell protein extraction buffer by Precellys 24 (Bertin Technologies, Paris, France) [23]. The cell homogenates were centrifuged at 10,800 g for 15 min at 4°C. The supernatant was used to perform RNA affinity purification.

Enrichment of RNA-binding proteins using RNA affinity purification

The rapid purification of RNA-binding protein streptavidin affinity tags was performed as described previously [12] with some modifications. Briefly, the streptavidin aptamer tagged RNAs were incubated with midgut cellular extract (pre-cleared with streptavidin beads) in



Figure 1. Identification of midgut cellular proteins that interact with the IFV 5' region of genomic RNA (A) Schematic of the secondary RNA structure generated for chromatography. The RNA construct contains the SP6 promoter fused to sequences from IFV sequence (nts 1–599) with the streptavidin aptamer inserted. (B) SDS-PAGE of protein that interacted with IFV 5' region of genomic RNA. Lane 'M' was prestained protein ladder, lane 'no RNA' was the supernatant that did not contain streptavidin aptamer-tagged RNA, lanes 'RNA' were RNA-binding proteins.

PBS containing 200 U of RNasin at 4°C for 2 h. RNA–protein complexes were affinity-purified with immobilized streptavidin beads. The RNA/protein/beads complexes were washed three times with cold PBS containing 200 U of RNasin. The elution was performed with saturated biotin solution and incubated at 4°C for 2 h. The beads without streptavidin aptamer-tagged RNA were used as the control. After centrifugation, the same volume of supernatant containing enriched proteins was resolved by 15% sodium dodecyl sulfatepolyacrylamide gel electrophoresis (SDS-PAGE). Supernatant without streptavidin aptamer-tagged RNA was used as a control and stored at -70°C until Q Exactive MS.

Mass spectrometry

The supernatant proteins (with and without aptamer-tagged RNA) were enriched three times, and pooled. The 150 µg protein of elution samples (RNA and no RNA system) was then digested using filteraided proteome preparation, lyophilized, and dissolved in 0.1% formic acid (FA). All of the peptide fractions were injected into the liquid chromatography (LC) system. The time of the LC gradient separation is 60 min. Experiments were performed on a Q Exactive mass spectrometer that was coupled to Easy nLC (Thermo Fisher Scientific, Waltham, USA). The precursor-ion mass tolerance was 20 ppm, and fragment-ion mass tolerance was 0.1 D. Each fraction (10 µl) was injected for nano-LC-MS/MS analysis. The peptide mixture (5 µg) was loaded onto a the C18-reversed phase column (Thermo Scientific Easy Column; 10 cm long, 75 µm inner diameter, and 3 µm resin) in buffer A (0.1% FA) and separated with a linear gradient of buffer B (80% acetonitrile and 0.1% FA) at a flow rate of 250 nl/min controlled by IntelliFlow technology over 60 min. MS data were acquired using a data-dependent top10 method dynamically choosing the most abundant precursor ions from the survey scan (300-1800 m/z) for higherenergy collisional dissociation (HCD) fragmentation. Determination of the target value is based on predictive automatic gain control. Dynamic exclusion duration was 40 s. Survey scans were acquired at a resolution of 70,000 at m/z 200 and resolution for HCD spectra was set to 17,500 at m/z 200. Normalized collision energy was 30 eV and the underfill ratio, which specifies the minimum percentage of the target value likely to be reached at maximum fill time, was defined as 0.1%. The instrument was run with peptide recognition mode enabled.

Western blot analysis

The supernatant proteins (with and without aptamer-tagged RNA) were separated by 12% SDS-PAGE and transferred onto nitrocellulose membranes (PALL, New York, USA) The membranes were blocked with 5% non-fat dried milk for 1 h at 37°C, and incubated with rabbit anti-coatomer subunit delta polyclonal antibodies (Abcam, Cambridge, UK), and rabbit anti-La protein homolog polyclonal antibodies which were prepared in our laboratory (data not shown). After washing, the membranes were incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG (Dingguo, Shanghai, China) and visualized by ECL Plus chemiluminescence reagent kit (Perkin-Elmer, Waltham, USA).

Data analysis

The raw MS/MS results were searched against the UniProt KB_ Bombyx mori database using Proteomics Tools (version 3.1.6) and Mascot (version 2.2). Database search parameters were as follows: enzyme was trypsin, missed cleavage was 2, static modifier was carbamidomethyl (C), dynamic modification was oxidation (M), and the filter parameters were protein FDR ≤ 0.01 and peptide FDR ≤ 0.01 .

The chromosome *loci* of the mapped genes were shown using Silk-Map tool (http://www.silkdb.org/silksoft/silkmap.html). The protein sequences were subject to BLAST query against the AmiGO database (http://amigo.geneontology.org/cgi-bin/amigo/blast.cgi) and clusters of orthologous groups (COGs) (http://www.ncbi.nlm.nih.gov/COG/). The corresponding gene ontology (GO) terms were extracted from the most homologous proteins, and the GO annotation results were plotted using the Web Gene Ontology Annotation Plotting (WEGO) (http://wego.genomics.org.cn/cgi-bin/wego/index.pl) by uploading compiled WEGO native format files containing the obtained GO terms [24].

The FASTA protein sequences of identified proteins were searched against KEGG (http://www.genome.jp/kegg/), and the corresponding KEGG pathways were extracted. Meanwhile, the interactional sites of the proteins of interest and IFV RNA were analyzed with catRAPID [25].

Results

RNA affinity capture of 5' region of IFV RNA-binding proteins

Silkworm midgut cellular proteins were identified that associate with an RNA containing sequences from the 5'-region of IFV and a

streptavidin RNA aptamer (Fig. 1A). Sequences from the 5'-region (nts 1–599) were amplified from IFV-CHN01 (GenBank: EU868609) [26]. In addition, the oligonucleotide used to amplify the 5' region incorporated a SP6 promoter for *in vitro* transcription. The obtained RNA-protein complexes were tested by SDS-PAGE. The results revealed that there were several weak stripes in systems with no RNA, while there were more bands in system with the 5' region of IFV RNA at the same volume supernatant (Fig. 1B).

Protein identification

In order to facilitate identification of proteins that associate with the IFV 5' region, the proteins were identified by PAGE-free MS. This methodology was applied to the interaction of RNA viruses and hosts, such as HCV, DENV, and FMDV [14,15,27] because it allows both identification of proteins in a complex mixture and direct comparison of recovered protein from the IFV 5'-region RNA sample. We found 4480 unique peptides and 726 protein groups (unique peptide ≥ 2) (Supplementary Table S1) by Q Exactive in RNA/protein/beads group with streptavidin aptamer-tagged RNA. The 325 protein groups were identified in system with RNA by comparing with system with no RNA (Supplementary Table S2).

Silkworms are female-heterogametic organisms (ZZ in male, ZW in female) that have 28 pairs of chromosomes [28]. To investigate the chromosome distribution of proteins that interacted with the 5'-region RNA of IFV and determine the authenticity of the proteins, 325 proteins were mapped onto the silkworm chromosomes using the corresponding CDS or amino acid sequence. Among 325 proteins, 311 were mapped onto the silkworm chromosome (**Supplementary Table S2**). All found proteins were relatively evenly distributed on all 28 chromosomes, with no obvious aggregation (**Fig. 2A** and **Supplementary Table S2**). The captured protein complexes were also tested by western blot analysis. Results showed that coatomer subunit delta and La protein homolog were detected by corresponding antibody (**Fig. 2B**).

*cat*RAPID was used to predict the interaction sites between eight identified proteins and the IFV 5'-region RNA. The region (from 100 to 350 nt) was the main binding region for these eight proteins (**Supplementary Fig. S1**). Based on *cat*RAPID fragments analysis, these proteins could bind to the 5' regions from 100 to 350 nt of IFV RNA, but the functions of these proteins remain to be verified. Human La protein could interact with GCAC near the initiator AUG enhance HCV RNA replication by promoting linkage between 5' and 3' UTR [29]. There was also GCAC (210–213 nt of IFV genome RNA) in the predicted RNA region that interacted with La protein homolog.

Function annotation

WEGO analysis classified the 325 proteins into three GO categories: cellular component (CC), molecular function (MF), and biological process (BP) (Fig. 3 and Supplementary Table S2). Under the CC category, large numbers of proteins were categorized as ribosome (GO:0005840), cytoplasm (GO:0005737), and integral to membrane (GO:0016021). IFV with other picornavirus proliferated in the cytoplasm, without entering into the nucleus [16], so these CCs were needed in life cycle of virus. ATP binding (GO:0005524), structural constituent of ribosome (GO:0003735), and nucleotide binding (GO:0000166) were the three most abundant subcategories under the MF category. Oxidation reduction (GO:0055114), proteolysis (GO:0006508), ribosome biogenesis (GO:0008152) accounted for



Figure 2. The confirmation of the identified proteins interacted with IFV 5' region of genomic RNA (A) Chromosome distribution of the identified proteins interacted with IFV 5' region of genomic RNA. (B) The western blot results of coatomer subunit delta and La protein homolog.



Figure 3. Gene ontology categories for the 325 proteins that interacted with IFV 5' region of genomic RNA The number of genes indicates those with available GO terms. The percentages indicate the proportion of the total genes.

the majority for the BP category. Picornavirus IRES-driven translation initiation depends on the recognition of the IRES by specific cellular proteins such as ATP, eIFs, and ribosome [8]. The COGs of proteins were generated by comparing the protein sequences of complete genomes [30]. Posttranslational modification, protein turnover, chaperones/translation, ribosomal structure, and

83

biogenesis/energy production and conversion were the top 3 identified functional classifications (**Supplementary Fig. S2** and **Table S3**).

KEGG pathway analysis

To provide a global view of functions of proteins that were identified, all proteins with different KO ids were submitted and mapped to 184 pathways (**Supplementary Table S4**). The top 24 pathways for identified proteins numbers are shown in Fig. 4. Metabolic pathway, biosynthesis, and microbial metabolism in diverse environments were the top three comprehensive pathways that were mapped. Ribosome and protein processing in endoplasmic reticulum were among the fourth and fifth pathways that were mapped (Figs. 4 and 5). These results further confirmed that 5' RNA region of IFV genome plays very important role in protein translation of IFV [18]. In addition, mRNA surveillance pathway (H9IYK6, H9JCI5, H9JG54, Q1HPL5, and H9JD63) and RNA transport (H9J5Q7, H9J7V0, H9JQ89, and Q684K3) were also found (**Supplementary Table S2** and S3). These proteins may play important roles in RNA process of IFV.

Discussion

The 5' region of IFV possesses an IRES [18], so the region might bind to translation-related factors and ribosomal proteins. In our results, 16 ribosomal proteins, 3 eIFs (eIF2, eIF3, and eIF4 subunit), elongation factor 1 (EF1), and six potential ITAFs were found to have RNA-binding motif (unique peptide \geq 2) (Table 1).

Initiation factors are necessary for processing mRNA prior to 40S subunit binding and the 60S subunit to form the 80S initiation complex [31]. However, different picornavirus IRESs need different eIFs [32]. For example, eIF4G, eIF4A, and eIF4B were shown to interact with type II IRES; however, eIF4E, eIF1, or eIF1A was not needed [33,34]. Poliovirus (PV) RNA utilizes eIF2 for the initiation of translation in cell-free system, and translation directed by PV RNA is blocked when eIF2 is inactivated at earlier time [35]. The two highly conserved RNA-binding motifs in eIF3 that direct translation initiation from the HCV IRES RNA [36]. The eIF4A subunits contain DEAD box domains, found within RNA helicases [37].

Elongation factors are a set of proteins in cellular protein synthesis. In the ribosome, they facilitate translational elongation, from the formation of the first peptide bond to the last one. Interactions between viral genomic RNA and eEF1 have been reported for several plant viruses, including tomato bushy stunt virus [38], tobacco mosaic virus [39], brome mosaic virus [40], turnip mosaic virus [41], turnip yellow mosaic virus [42], and animal viruses including bovine viral diarrhea virus [43], PV [44], West Nile virus [45,46], and Dengue 4 virus [47]. Elongation factors play important roles in viral translation and replication.

Several potential picornavirus ITAFs were found in this study, such as La (Table 1). The ITAFs are important *trans*-acting factors required for viral translation and replication [10]. La could enhance IRESmediated translation in PV, EMCV, and HCV [48–50]. The importance of nucleoplasmin family in cellular processes such as chromatin remodeling, genome stability, ribosome biogenesis, DNA duplication, and transcriptional regulation has led to the rapid growth of information available on their structure and function [51]. A conserved RNA recognition motif protein encoded by *LARK* gene is eliminated to determine the consequences for circadian timing *in vivo* [52] that might affect the IFV replication and host life cycle.

The cytoskeleton and motor proteins have been found to play important roles in RNA intracellular localization and may also regulate translation and viral replication [53–55]. Twelve cytoskeleton and proteins were identified as tubulin and myosin etc. (Table 2). Riboproteomic analysis of polypeptides interacting with the IRES of foot-and-mouth disease virus and HCV RNA revealed cytoskeleton-related proteins [14,15]. Myosin plays a key role in the transport of cargo to the plasma membrane. Its cargo includes cell-surface receptors, pigment, and organelles such as the endoplasmic reticulum. It is also emerging that RNA and RNA–BPs (RNA-binding proteins) make up another class of myosin cargo [56].

Membrane lipid composition is important for RNA replication of positive-strand RNA viruses [57–60]. Several kinds of membrane-related proteins were found as shown in Table 3. The membrane-shaping host reticulon proteins play crucial roles in viral RNA replication compartment formation and function [61,62]. With IFV infected silkworm, goblet cell cytoplasm was filled with specific membranous vesicles and



Figure 4. Top 24 KEGG pathways for identified protein genes





Figure 5. Distribution of the 16 identified ribosome subunits and 15 proteins in protein processing in endoplasmic reticulum pathway interacted with IFV 5' region of genomic RNA Red labeled proteins were identified by this study.

Accession number	Protein description	PepCount	Unique PepCount	Cover percent	MW (Da)	PI	Length
Q5UAL6	40S ribosomal protein S27	3	2	27.38	9433.04	9.91	84
Q5UAN5	Ribosomal protein S9	4	4	19.07	22,576.11	10.76	194
Q6PUF7	40S ribosomal protein S24	4	2	23.48	15,171.86	10.91	132
Q5UAM7	Ribosomal protein S15A	2	2	13.18	14,692.17	10.01	129
Q5UAS7	Ribosomal protein L12	3	3	19.51	17,565.2	9.58	164
Q5UAM2	Ribosomal protein S20	4	2	19.51	13,731.99	10.14	123
Q5UAR8	Ribosomal protein L21	2	2	12.58	17,984.96	10.96	159
Q5UAQ1	Ribosomal protein L36A	2	2	20.19	12,294.53	10.53	104
Q5UAP9	Ribosomal protein L37A	2	2	16.67	10,031.87	10.6	90
Q5UAQ8	Ribosomal protein L30	6	5	35.40	12,346.36	9.65	113
H9J1M5	Ribosomal protein LP1	6	4	56.25	11,478.77	4.18	112
H9IVT2	40S ribosomal protein S21	2	2	23.46	8811.06	7.67	81
Q5UAN7	Ribosomal protein S7	6	2	18.95	21,858.28	9.82	190
H9JX29	Ribosomal protein L14	7	3	20.61	18,623.75	11.27	165
Q5UAM5	Ribosomal protein S17	5	4	42.11	15,421.65	9.71	133
H9JU64	Ribosomal protein L6	8	6	20.38	28,999.88	10.89	260
Q684K3	Translation initiation factor 2 gamma	5	4	10.90	50,600.37	9.03	468
H9JTH1	Eukaryotic translation initiation factor	2	2	11.03	31,411.22	9.76	290
H9J7V0	Eukaryotic translation initiation factor 3	2	2	2.91	62,410.69	7.58	549
H9JCI5	EF1 alpha-like factor isoform 1	2	2	6.97	65,964.74	5.16	603
Q2F5N9	Nucleoplasmin isoform 2	3	2	22.99	20,843.06	4.86	187
H9JJU3	Aspartyl-tRNA synthetase	2	2	4.36	46,355.27	9.03	413
H9J930	La protein homolog	2	2	8.53	45,133.49	6.01	387
Q5I7M6	RNA-binding protein lark	5	5	15.74	38,515.19	9.31	343
Q17201	Bmsqd-2	4	2	9.24	32,552.06	8.66	303

Table 1. Identification of translation initiation, elongation, and potential IRES trans-acting factors that interact with IFV 5'-region RNA

Table 2. Identification of cytoskeleton and motor-related that interact with IFV 5'-region RNA

Accession number	Protein description	PepCount	Unique PepCount	Cover percent	MW (Da)	PI	Length
H9J4V2	Tubulin beta-4b chain	9	4	13.86	37,243.67	4.91	332
Н9ЈНХЗ	Isoform F	3	3	1.85	209,582.44	6.49	1943
H9J4H4	Short isoform H	2	2	1.37	203,746.09	5.72	1829
H9J1Q5	Tubulin beta-4b chain-like isoform x1	15	7	20.96	49,056.33	5.79	439
H9J3F1	Microtubule-associated protein	2	2	8.57	39,211.21	5.74	350
H9JGR0	Actin-interacting protein 1-like	3	2	18.63	17,422.34	7.93	161
B3VTP0	Long form-like	18	16	21.41	102,749.53	5.43	878
H9JXG0	Myosin heavy muscle	26	23	18.98	134,836.77	5.97	1175
H9JXG1	Muscle myosin heavy chain	78	53	53.33	123,511.13	5.36	1082
B5M9A1	Myosin heavy chain	37	28	36.54	91,159.53	6.54	791
C4P7H2	Myosin heavy chain variant A	17	15	25.43	65,828.31	6.24	582

electron-dense bodies [16]. PV replicates its genome in association with membranous vesicles in the cytoplasm of infected cells, and cellular COPII coat proteins are involved in production of the vesicles that form the PV replication complex [63–65]. Although we did not found COPII coat proteins, three subunits of coatomer protein that own COPI vesicle coat (GO:0030126) and retrograde vesicle-mediated transport, Golgi to endoplasmic reticulum (GO:0006890) were found (Table 3). In addition, other found membrane-related proteins (Table 3) could better explore the mechanism of membranous vesicles forming in the cytoplasm. The forming mechanism of specific vesicles and electron-dense bodies may be related to IFV 5'-region RNA.

Interestingly, we identified a number of enzymes involved in ATP, mitochondrial-related and metabolic enzymes (**Supplementary Tables S5** and **S6**). Mitochondrial ATP synthase is a factor that mediates HIV-1 transfer between APCs and CD4(+) target cells [66].

Purified mitochondria from HIV infected cells are infectious, and mitochondrial inhibitors block HIV transmission [67]. After silkworms are infected with IFV, a number of mitochondria surround the virions [16]. Based on this, we hypothesize that ATP synthase and mitochondrial-related proteins play important roles in viral transmission and life cycle. Picornavirus IRES-driven translation initiation depends on the recognition of the IRES by ATP [8].

Metabolic enzymes that bind RNA form another level of the cellular regulatory network [68]. However, the function of nucleotidebinding activity by carbohydrate metabolic enzyme is unclear. RNAbinding activity has been reported, such as for alpha-glucosidase and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) [69]. The NAD⁺-binding Rossman domain found within GAPDH and other dehydrogenase- and nucleotide-binding family proteins should be designated as a general RNA-binding motif [69,70]. Table 3. Identification of membrane-related that interact with IFV 5'-region RNA

Accession number	Protein description	PepCount	Unique PepCount	Cover percent	MW (Da)	PI	Length
Q1HQA9	Transmembrane trafficking protein	3	2	12.20	23,681.24	6.01	205
Q14UC7	Aquaporin AQP-Bom2	2	2	10.42	27,693.23	6.17	259
Q6SA04	Innexin 2	3	3	12.81	41,810.52	6.6	359
H9IT08	Leucine-rich repeat-containing protein 15-like	7	6	3.63	232,055.3	4.73	2095
H9JKN7	AFG3-like protein 2-like	2	2	2.85	85,489.25	8.75	773
H9IYQ9	Calcium-binding mitochondrial carrier protein	3	3	8.85	74,423.77	8.88	667
Н9ЈМН2	Coatomer subunit alpha	4	4	4.10	131,334.2	6.97	1170
H9JSC6	Coatomer subunit epsilon	2	2	11.59	33,251.22	5.09	302
H9IS78	Coatomer subunit delta	4	3	5.43	63,528.16	6.56	571
H9JI54	Organic cation transporter	2	2	7.88	60,897.33	6.56	546
Q1HPK0	Synaptic vesicle membrane protein vat-1	2	2	7.14	48,819.91	5.85	448
Q2F5Y0	NADH dehydrogenase	2	2	9.95	25,036.05	6.15	221
H9J1D9	Membrane protein precursor	3	3	12.95	24,251.66	4.62	224
H9J1D5	Membrane protein precursor	4	4	29.20	24,824.91	4.7	226
H9JXN1	Membrane protein precursor	5	4	15.75	27,854.72	5	254
H9J971	Sodium potassium-transporting atpase	2	2	3.08	63,324.45	7.59	584
H9IUU2	CD9 antigen-like	3	2	21.05	20,872.39	4.83	190
H9J9I6	dipeptidyl peptidase 4	4	3	7.29	80,264.2	5.09	713
H9IXU1	Anion exchange protein slc4a2	2	2	3.52	93,447.19	8.26	853
H9JL65	Low quality protein: de-cadherin	2	2	3.72	90,520.7	4.57	807
H9JFQ1	Niemann-pick c1	2	2	1.18	247,266.8	4.81	2206
H9JAG3	Transmembrane emp24 protein 7	4	2	28.82	19,472.75	5.6	170
H9J4U3	Plasma membrane calcium-transporting atpase	2	2	10.30	36,860.91	6.72	330
H9IVS2	Mannose binding	2	2	6.43	31,252.26	6	280
H9JRC2	Isoform A	2	2	7.11	49,871.97	6.6	464
H9IUL3	Synaptobrevin-like isoform 1	3	2	20.00	13,136.01	9.05	120
Q5CCJ5	Ras-like protein 2-like	2	2	10.50	22,879.86	6.34	200



Figure 6. Summary of IFV 5'-end RNA interacting proteins identified by MS These interactions will require further study to examine their function, involved links

between the IFV 5'-end RNA and the cytoskeleton, metabolism, and translation.

Cellular chaperones and folding enzymes are vital contributors to membrane-bound replication and movement complexes during RNA virus infection [71-73]. The folding enzymes (Supplementary Table S7) and GTP-binding proteins (Supplementary Table S8) were also found. The small GTP-binding protein ARL8 plays crucial roles in tobamovirus RNA replication [74]. Histone deacetylase activity may be required for the virus replication [75]. Specifically, the histones H2B and H4 were identified. The phosphorylation of viral and cellular proteins can have major impacts on viral infection, replication, and cytotoxicity in the host cell [76]. Three serine/threonine-protein phosphatases (Q1HPL5, H9JG54, and H9JD63) were found. The replication of positive-strand RNA viruses, the largest class of viruses, including significant human, animal, and plant pathogens, may also be regulated by phosphorylation events [77,78]. In addition, 30 KD proteins (30KPs) (Supplementary Table S9) may play an important role in translation and replication of IFV. The 30KPs were the major plasma proteins and classified into the lepidopteran-specific lipoprotein 11 family [79]. The 30KPs in silkworm may serve as the amino acid source during embryo development and play a role in the inhibition of apoptosis [79,80].

This study aimed to identify proteins that interacted with host factors and the 5' end of IFV genomic RNA. The identification of many proteins (ribosome, endoplasmic reticulum, membrane, mitochondrial, and IRES-related proteins) with potential roles in viral translation and replication indicates that much work is needed to better understand these complex functions and pathways (Fig. 6). These results also revealed many enzymes that may serve multiple roles during IFV infection of the silkworm midgut. Functional investigations of identified RNA-binding proteins of interest are needed to determine their role in viral translation and replication, and to provide potential therapeutic targets.

Supplementary Data

Supplementary Data is available at ABBS online.

Funding

This work was supported by the grants from National Natural Science Foundation of China (No. 31172262) and the Postdoctoral Foundation of China (No. 2013M531471).

References

- Lu HS. Molecular Biology of Insect Viruses. Beijing: Chinese Agricultural Science Press, 1998.
- 2. van Oers MM. Insect Virology. Norfolk: Caister Academic Press, 2010.
- Liu Y, Wimmer E, Paul AV. cis-acting RNA elements in human and animal plus-strand RNA viruses. Biochim Biophys Acta 2009, 1789: 495–517.
- Martinez-Salas E. The impact of RNA structure on picornavirus IRES activity. Trend Microbiol 2008, 16: 230–237.
- Lukavsky PJ. Structure and function of HCV IRES domains. Virus Res 2009, 139: 166–171.
- Nagy PD, Pogany J. The dependence of viral RNA replication onco-opted host factors. *Nat Rev Microbiol* 2012, 10: 137–149.
- Li Z, Nagy PD. Diverse roles of host RNA binding proteins in RNA virus replication. RNA Biol 2011, 8: 305–315.
- Martinez-Salas E, Pacheco A, Serrano P, Fernandez N. New insights into internal ribosome entry site elements relevant for viral gene expression. *J Gen Virol* 2008, 89: 611–626.

- Fernandez-Miragall O, Lopez de Quinto S, Martinez-Salas E. Relevance of RNA structure for the activity of picornavirus IRES elements. *Virus Res* 2009, 139: 172–182.
- Pacheco A, Martinez-Salas E. Insight into the biology of IRES element through riboproteomic approaches. J Biomedic and Biotech 2010, 2010: 458927.
- Schmidt C, Kramer K, Urlaub H. Investigation of protein-RNA interactions by mass spectrometry-techniques and applications. *J Proteomics* 2012, 75: 3478–3494.
- Lu H, Li WQ, Noble WS, Payan D, Anderson DC. Riboproteomics of the hepatitis C virus internal ribosomal entry site. J Proteome Res 2004, 3: 949–957.
- Pacheco A, Reigadas S, Martinez-Salas E. Riboproteomic analysis of polypeptides interacting with the internal ribosome-entry site element of foot-and-mouth disease viral RNA. *Proteomics* 2008, 8: 4782–4790.
- Walker SC, Scott FH, Srisawat C, Engelke DR. RNA affinity tags for the rapid purification and investigation of RNAs and RNA-protein complexes. *Methods Mol Biol* 2008, 488: 23–40.
- Sawicka K, Bushell M, Spriggs KA, Willis AE. Polypyrimidine-tract-binding protein: a multifunctional RNA-binding protein. *Biochem Soc Trans* 2008, 36: 641–647.
- Man NN, Lu QN, Hong J, Jin W, Lu XM. Immunoelectron micrography of the midgut epithelium infected with *Bombyx mori* infectious flacherie virus. *Sci Seric* 2007, 33: 602–609.
- Hashimoto Y, Watanabe A, Kawase S. *In vitro* translation of infectious flacherie virus RNA in a wheat germ and a rabbit reticulocyte system. *Biochim Biophys Acta* 1984, 781: 76–80.
- Li MQ, Man NN, Qiu HH, Cai SF, He XY, He XK, Lu XM. Detection of an internal translation activity in the 5' region of *Bombyx mori* infectious flacherie virus. *Appl Microbiol Biotechnol* 2012, 95: 697–705.
- Ongus JR, Roode EC, Pleij CW, Vlak JM, van Oers MM. The 5' nontranslationed region of Varroa destructor virus (genus iflavirus): structure prediction and IRES activity in *Lymantria dispar* cells. J Gen Virol 2006, 87: 3397–3407.
- Wu TY, Wu CY, Chen YJ, Chen CY, Wang CH. The 5' untranslated region of *Perina nuda* virus (PnV) possesses a strong internal translation activity in baculovirus-infected insect cells. *FEBS Lett* 2007, 581: 3120–3126.
- 21. Lu J, Zhang J, Wang X, Jiang H, Liu C, Hu YY. *In vitro* and *in vivo* identification of structural and sequence elements in the 5' untranslated region of Ectropis oblique picornalike virus required for internal initiation. *J Gen Virol* 2006, 87: 3667–3677.
- Srisawat C, Engelke DR. Streptavidin aptamers: affinitytags for the study of RNAs and ribonucleoproteins. RNA 2001, 7: 632–641.
- Wu H, Southam AD, Hines A, Viant MR. High-throughput tissue extraction protocol for NMR- and MS-based metabolomics. *Anal Biochem* 2008, 372: 204–212.
- 24. Ye J, Fang L, Zheng H, Zhang Y, Chen J, Zhang ZJ, Wang J, et al. WEGO: a web tool for plotting GO annotations. Nucleic Acids Res 2006, 34: W293–W297.
- Bellucci M, Agostini F, Masin M, Tartaglia GG. Predicting protein associations with long noncoding RNAs. Nat Methods 2011, 8: 444–445.
- 26. Li MQ, Chen XX, Wu XX, Man NN, Jin W, Lu XM. Genome analysis of the *Bombyx mori* infectious flacherie virus isolated in China. Agr Sci in China 2010, 9: 299–305.
- 27. Ward AM, Bidet K, Yinglin A, Ler SG, Hogue K, Blackstock W, Gunaratne J, et al. Quantitative mass spectrometry of DENV-2 RNA-interacting proteins reveals that the DEAD-box RNA helicase DDX6 binds the DB1 and DB2 3' UTR structures. RNA Biol 2011, 8: 1173–1186.
- Hasimoto H. The role of the W-chromosome in the sex determination of Bombyx mori. Jpn J Genet 1993, 8: 245–247.
- 29. Kumar A, Ray U, Das S. Human La protein interaction with GCAC near the initiator AUG enhances hepatitis C Virus RNA replication by promoting linkage between 5' and 3' untranslated regions. *J Virol* 2013, 87: 6713–6726.
- Tatusov RL, Galperin MY, Natale DA, Koonin EV. The COG database: a tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res* 2000, 28: 33–36.

- Pain VM. Initiation of protein synthesis in eukaryotic cells. Eur J Biochem 1996, 236: 747–771.
- Jackson RJ, Hellen CU, Pestova TV. The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat Rev Mol Cell Biol* 2010, 11: 113–127.
- 33. Kolupaeva VG, Lomakin IB, Pestova TV, Hellen CU. Eukaryotic initiation factors 4G and 4A mediate conformational changes downstream of the initiation codon of the encephalomyocarditis virus internal ribosomal entry site. *Mol Cell Biol* 2003, 23: 687–698.
- 34. Kolupaeva VG, Pestova TV, Hellen CU, Shatsky IN. Translation eukaryotic initiation factor 4G recognizes a specific structural element within the internal ribosome entry site of encephalomyocarditis virus RNA. J Biol Chem 1998, 273: 18599–18604.
- Redondo N, Sanz MA, Welnowska E, Carrasco L. Translation without eIF2 promoted by poliovirus 2A protease. *PLoS One* 2011, 6: e25699.
- 36. Sun C, Querol-Audí J, Mortimer SA, Arias-Palomo E, Doudna JA, Nogales E, Cate JH. Two RNA-binding motifs in eIF3 direct HCV IRESdependent translation. *Nucleic Acids Res* 2013, 41: 7512–7521.
- Rogers GW, Komar AA, Merrick WC. eIF4A: the godfather of the DEAD box helicases. Prog Nucleic Acid Res Mol Biol 2002, 72: 307–331.
- Li ZH, Pogany J, Tupman S, Esposito AM, Kinzy TG, Nagy PD. Translation elongation factor 1A facilitates the assembly of the tombusvirus replicase and stimulates minus-strand synthesis. *PLoS Pathog* 2010, 6: e1001175.
- Hwang J, Oh CS, Kang BC. Translation elongation factor 1B (eEF1B) is an essential host factor for Tobacco mosaic virus infection in plants. *Virology* 2013, 439: 105–114.
- Bastin M, Hall TC. Interaction of elongation-factor 1 with aminoacylated Brome mosaic virus and transfer RNAs. J Virol 1976, 20: 117–122.
- 41. Thivierge K, Cotton S, Dufresne PJ, Mathieu I, Beauchemin C, Ide C, Fortin MG, *et al*. Eukaryotic elongation factor 1A interacts with turnip mosaic virus RNA-dependent RNA polymerase and VPg-Pro in virus-induced vesicles. *Virology* 2008, 377: 216–225.
- Joshi RL, Ravel JM, Haenni AL. Interaction of turnip yellow mosaic virus Val-RNA with eukaryotic elongation factor EF-1 (alpha]. Search for a function. *EMBO J* 1986, 5: 1143–1148.
- 43. Johnson CM, Perez DR, French R, Merrick WC, Donis RO. The NS5A protein of bovine viral diarrhoea virus interacts with the alpha subunit of translation elongation factor-1. J Gen Virol 2001, 82: 2935–2943.
- 44. Harris KS, Xiang W, Alexander L, Lane WS, Paul AV, Wimmer E. Interaction of poliovirus polypeptide 3CDpro with the 5' and 3' termini of the poliovirus genome. Identification of viral and cellular cofactors needed for efficient binding. J Biol Chem 1994, 269: 27004–27014.
- 45. Blackwell JL, Brinton MA. Translation elongation factor-1 alpha interacts with the 3' stem-loop region of West Nile virus genomic RNA. J Virol 1997, 71: 6433–6444.
- 46. Davis WG, Blackwell JL, Shi PY, Brinton MA. Interaction between the cellular protein eEF1A and the 30-terminal stem-loop of West Nile virus genomic RNA facilitates viral minus-strand RNA synthesis. J Virol 2007, 81: 10172–10187.
- 47. De Nova-Ocampo M, Villegas-Sepuveda N, del Angel RM. Translation elongation factor-1 alpha, La, and PTB interact with the 3' untranslated region of dengue 4 virus RNA. *Virology* 2002, 295: 337–347.
- Kim YK, Jang SK. La protein is required for efficient translation driven by encephalomyocarditis virus internal ribosomal entry site. J Gen Virol 1999, 80: 3159–3166.
- 49. Izumi RE, Das S, Barat B, Raychaudhuri S, Dasgupta A. A peptide from autoantigen La blocks poliovirus and hepatitis C virus cap-independent translation and reveals a single tyrosine critical for La RNA binding and translation stimulation. J Virol 2004, 78: 3763–3776.
- Meerovich K, Svitkin YV, Lee HS, Lejbkowicz F, Kenan DJ, Chan EK, Agol VI, et al. La autoantigen enhances and corrects aberrant translation of poliovirus RNA in reticulocyte lysate. J Virol 1993, 67: 3798–3807.
- Frehlick LJ, Eirín-López JM, Ausió J. New insights into the nucleophosmin/ nucleoplasmin family of nuclear chaperones. *Bioessays* 2007, 29: 49–59.
- 52. Sundram V, Ng FS, Roberts MA, Millán C, Ewer J, Jackson FR. Cellular requirements for LARK in the *Drosophila circadian* system. J Biol Rhythms 2012, 27: 183–195.

- Lopez de Heredia M, Jansen RP. mRNA localization and the cytoskeleton. Curr Opin Cell Biol 2004, 16: 80–85.
- Bost A, Venable D, Liu L, Heinz B. Cytoskeletal requirements for hepatitis C virus (HCV) RNA synthesis in the HCV replicon cell culture system. *J Virol* 2003, 77: 4401–4408.
- Dohner K, Sodeik B. The role of the cytoskeleton during viral infection. *Curr Top Microbiol Immunol* 2005, 285: 67–108.
- McCaffrey MW, Lindsay AJ. Roles for myosin Va in RNA transport and turnover. *Biochem Soc Trans* 2012, 40: 1416–1420.
- 57. Lee WM, Ishikawa M, Ahlquist P. Mutation of host delta 9 fatty acid desaturase inhibits brome mosaic virus RNA replication between template recognition and RNA synthesis. J Virol 2001, 75: 2097–2106.
- Sharma M, Sasvari Z, Nagy PD. Inhibition of sterol biosynthesis reduces tombusvirus replication in yeast and plants. J Virol 2010, 84: 2270–2281.
- Mackenzie JM, Khromykh AA, Parton RG. Cholesterol manipulation by West Nile virus perturbs the cellular immune response. *Cell Host Microbe* 2007, 2: 229–239.
- Ahola T, Lampio A, Auvinen P, Kaariainen L. Semliki forest virus mRNA capping enzyme requires association with anionic membrane phospholipids for activity. *EMBO J* 1999, 18: 3164–3172.
- Diaz A, Wang X, Ahlquist P. Membrane-shaping host reticulon proteins play crucial roles in viral RNA replication compartment formation and function. *PNAS* 2010, 107: 16291–16296.
- Salonen A, Ahola T, Kääriäinen L. Viral RNA replication in association with cellular membranes. *Curr Top Microbiol Immunol* 2005, 285: 139–173.
- Rust RC, Landmann L, Gosert R, Tang BL, Hong W, Hauri HP, Egger D, et al. Cellular COPII proteins are involved in production of the vesicles that form the poliovirus replication complex. J Virol 2001, 75: 9808–9818.
- Barco A, Carrasco L. A human virus protein, poliovirus 2BC, induces membrane proliferation and blocks the exocytic pathway in the yeast Sacchromyces cere-6isiae. EMBO J 1995, 14: 3349–3364.
- Schlegel A, Giddings TH Jr, Ladinsky MS, Kirkegaard K. Cellular origin and ultrastructure of membranes induced during poliovirus infection. J Virol 1996, 70: 6576–6588.
- 66. Yavlovich A, Viard M, Zhou M, Veenstra TD, Wang JM, Gong W, Heldman E, *et al.* Ectopic ATP synthase facilitates transfer of HIV-1 from antigen-presenting cells to CD4+ target cells. *Blood* 2012, 120: 1246–1253.
- Lee-Huang S, Huang PL, Lee-Huang P. Live-cell real-time imaging reveals role of mitochondria in cell-to-cell transmission of HIV-1. *Biochem Biophys Res Comm* 2011, 415: 384–389.
- Ciesla J. Metabolic enzymes that bind RNA: yet another level of cellular regulatory network. *Acta Biochim Pol* 2006, 53: 11–32.
- Doroshenk KA, Crofts AJ, Morris RT, Wyrick JJ, Okita TW. Proteomic analysis of cytoskeleton-associated RNA binding proteins in developing rice seed. J Proteome Res 2009, 8: 4641–4653.
- Nagy E, Henics T, Eckert M, Miseta A, Lightowlers RN, Kellermayer M. Identification of the NAD+-binding fold of glyceraldehyde-3-phosphate dehydrogenase as a novel RNA-binding domain. *Biochem Biophys Res Commun* 2009, 275: 252–260.
- Verchot J. Cellular chaperones and folding enzymes are vital contributors to membrane bound replication and movement complexes during plant RNA virus infection. *Front Plant Sci* 2012, 3: 275.
- Nagy PD, Wang RY, Pogany J, Hafren A, Makinen K. Emerging picture of host chaperone and cyclophilin roles in RNA virus replication. *Virology* 2011, 411: 374–382.
- Zúñiga S, Sola I, Cruz JL, Enjuanes L. Role of RNA chaperones in virus replication. Virus Res 2009, 139: 253–266.
- 74. Nishikiori M, Mori M, Dohi K, Okamura H, Katoh E, Naito S, Meshi T, et al. A host small GTP-binding protein ARL8 plays crucial roles in tobamovirus RNA replication. PLoS Pathog 2011, 7: e1002409.
- 75. Shim SH, Park JH, Ye MB, Nam JH. Histone deacetylase inhibitors suppress coxsackievirus B3 growth *in vitro* and myocarditis induced in mice. *Acta Virol* 2013, 57: 462–466.
- Keatin JA, Striker R. Phosphorylation events during viral infections provide potential therapeutic targets. *Rev Med Virol* 2012, 22: 166–181.

- 77. Jakubiec A, Jupin I. Regulation of positive-strand RNA virus replication: the emerging role of phosphorylation. *Virus Res* 2007, 129: 73–79.
- Zhang Y, Dong Z, Liu S, Yang Q, Zhao P, Xia QY. Identification of novel members reveals the structural and functional divergence of lepidopteranspecific Lipoprotein_11 family. *Funct Integr Genomics* 2012, 12: 705–715.
- 79. Zhong BX, Li JK, Lin JR, Liang JS, Su SK, Xu HS, Yan HY, et al. Possible effect of 30 K proteins in embryonic development of silkworm *Bombyx* mori. Acta Biochim Biophys Sin 2005, 37: 355–361.
- Kim EJ, Park HJ, Park TH. Inhibition of apoptosis by recombinant 30 K protein originating from silkworm hemolymph. *Biochem Biophys Res Commun* 2003, 308: 523–528.