

Requirement of crude protein for maintenance in a new strain of laboratory rabbit

Jian-Min Lv^{a,b}, Min-li Chen^{b,*}, Li-Chun Qian^a, Hua-Zhong Ying^b, Jian-Xin Liu^{a,**}

^a College of Animal Sciences, Zhejiang University, Huajiachi Campus, Hangzhou 310029, China

^b Laboratory Animal Research Center, Zhejiang Chinese Medicine University, Hangzhou 310053, China

ARTICLE INFO

Article history: Received 13 April 2008 Received in revised form 30 October 2008 Accepted 24 December 2008

Keywords: Crude protein Requirement for maintenance Nitrogen balance Rabbit

ABSTRACT

A new strain of laboratory rabbit, produced from Japanese White rabbit but characterized by its black eye and higher antibody titer, was used in a series of nitrogen (N) balance trials to investigate the requirement of crude protein (CP) for maintenance. The N balance was determined for growing (6-12 weeks) and finishing period (13-17 weeks). Thirty rabbits were used in each period. The rabbits in both periods were divided into five equal groups and fed the diets with equal digestible energy but different contents of CP (120, 140, 160, 180 and 200 g/kg DM). The N requirements for maintenance were estimated from the relationship between the N retention and N intake. Following the above series of N balance tests, 20 rabbits of 16 of age weeks were used with a N-free diet. The rabbits were divided into five groups and offered the N-free diet at levels of 55, 45, 35, 25 and 0 g/day, respectively. Net protein utilization ranged from 0.45 to 0.50 and increased with the advancing age of rabbits, but no significant difference was found between different contents of dietary CP. The estimated N requirement for maintenance was, on average, 485 mg/kg BW^{0.75} per day, equivalent to 3.03 g CP/kg BW^{0.75}. The result from the trial with the N-free diet showed that N requirement for maintenance was 486 mg/kgBW^{0.75} per day, confirming the results obtained in the series of N balance tests. The

0377-8401/\$ – see front matter © 2009 Elsevier B.V. All rights reserved.

doi:10.1016/j.anifeedsci.2009.01.001

Abbreviations: N, nitrogen; CP, crude protein; WHBE, white hair and black eye; NPU, net protein utilization; NI, N intake; NR, N retention; Nm, nitrogen for maintenance; TNE, total nitrogen excreted in feces and urine.

^{*} Corresponding author. Tel.: +86 571 86613590; fax: +86 571 86613662.

^{**} Corresponding author. Tel.: +86 571 86971097; fax: +86 571 86971930. E-mail addresses: cml991@yahoo.com.cn (M.-l. Chen), liujx@zju.edu.cn (J.-X. Liu).

lower intake of the N-free diet resulted in more N excretion suggesting that protein catabolism may occur in the body of rabbit to meet maintenance requirements for N when the dietary N intake was very low.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

The domestic rabbit (*Oryctolagus cuniculus*) was descended from the European wild rabbits and developed into various breeds of modern domestic rabbits from the 18th century. There are now several hundred varieties throughout the world varying in size, colour, type of hair coat and other characteristics (NRC, 1977). These breeds could be used as different purposes such as for meat, wool and for research. The use of laboratory animals is considered to be necessary, for testing new surgical techniques, the study of new chemical and pharmaceutical substances and the production of vaccines and antibodies (Amalia, 2004). The preparation of antiserum is one of the most important usages of laboratory rabbits (Wei, 1998).

A new strain of laboratory rabbit with consistently inherited black eye trait was produced by inbreeding from a mutant of the Japanese White rabbit. Different to the red eye color of the Japanese White rabbit, this new strain is characterized by black eye trait, so it has been known as white hair and black eye (WHBE). It has been observed that the WHBE rabbit was sensitive to changes in environment and adapted for biomedical research, especially in immunology research. Compared with other breeds of laboratory rabbits, the WHBE rabbit could produce higher antibody titer response to some special antigen (Chen et al., 2005). As far as these special characteristics in WHBE rabbits were concerned, it was assumed that their nutritional requirements may be different from other breeds. Amalia (2004) suggested that laboratory rabbits had quite different nutrient requirements from rabbits that were sold commercially. However, due to the absence of feed standards for laboratory rabbit, researchers have to rely on the nutrient requirements of meat-type rabbits or personal experience in determining the feed allowances, which may compromise the ability of the WHBE rabbit to express maximal potential in immune and other biological aspects. Furthermore, excess dietary nitrogen could not only place strain on the kidney, but also cause environmental pollution (Maerten et al., 1997). It is, therefore, necessary to establish feeding allowances for WHBE rabbits.

Energy and protein (amino acids) are the most critical dietary components for animals. Although the requirements of protein (amino acids) for meat-type rabbits have been established (NRC, 1977), little information is available for laboratory rabbits. Most studies of the nutrient requirement of rabbits were based on feeding trials either in China or abroad (Li and Wei, 2002; NRC, 1977; Maerten, 1992) and there was little examination of nutritional parameters. In factorial analysis, the requirements for nutrients may be divided into various components including maintenance, weight gain, hair growth, *etc.* With respect to protein (amino acid) requirements, that for maintenance is significant and it is important that it is quantified such that total requirements can be established allowing differences between animal species to be determined.

In the current study, a series of nitrogen (N) balance trials were carried out to investigate the requirement of crude protein (CP) for maintenance in WHBE rabbits.

2. Materials and methods

These experiments were approved by the Animal Care and Use Committee of Zhejiang Province and carried out in Laboratory Animal Research Center, Zhejiang Chinese Medicine University, China. The rabbits used in the trials were in good health and met the criteria for microbiology and parasitology (AQSIQ, 2001a,b). The numbers of laboratory-animal-quality certification and experimental facility license were SCXK(ZHE) 2005-0022 and SYXK(ZHE) 2003–0004, respectively.

Items ^a	Crude protein (g/kg)						
	120	140	160	180	200		
Digestible energy (MJ/kg DM)	10.4	10.4	10.4	10.4	10.4		
Crude protein (g/kg DM)	115	138	159	175	205		
Crude fibre (g/kg DM)	123	125	113	115	116		
Ca (g/kg DM)	15	15	15	15	15		
P(g/kg DM)	8	8	8	8	8		
Lysine (g/kg DM)	6.6	7.9	9.0	10.1	11.3		
Methionine + cysteine (g/kg DM)	4.9	5.7	6.5	7.3	8.1		
Threonine (g/kg DM)	5.0	5.8	6.7	7.5	8.3		

The composition of experimental diets used in the series of nitrogen balance tests.

^a The digestible energy, Ca, P and amino acids were calculated from the ingredients of the diets, while the contents of crude protein and crude fibre were determined values (AOAC, 1990).

2.1. Nitrogen balance test

Table 1

Sixty weanling rabbits (42-day-age, male:female = 1:1) were selected and housed individually in cages in a room with controlled temperature and a regular photoperiod (12 h light and 12 h dark) and given free access to feed and water. Nitrogen balance tests were carried out at the age of 10 and 15 weeks, corresponding to the growing and finishing periods, respectively. In each period, 30 rabbits (male:female = 1:1) were selected and divided into 5 equal groups. Each group with six replicates was randomly allocated to an isoenergetic ration with different content of CP ranging from 120 to 200 g/kg (Table 1). In all experimental diets, the contents of lysine, methionine plus cystine and threonine in CP were similar. All the rabbits for N balance test were confined in individual metabolism cages for collection of faeces and the data were individually recorded.

Each period consisted of 12 days, the first 9 days of which were for adaptation and the subsequent 3 days for collection. During the latter period, faeces and urine were collected respectively at 8:00 a.m. every day. Total faecal output for 3 days was mixed with 20 ml tartaric acid (10%) and dried in an air-force drier at 65–70 °C. After being mixed, ground (40 Mesh) and weighed, a 10% sample of faeces was retained. Urine was acidified with 10% tartaric acid and refrigerated at 4 °C until the end of the collection period. A 10% sample of urine was retained and frozen for subsequent N analysis. Feed, faeces and urine samples were analyzed for N content by Kjeldahl process (methods number 984.13; AOAC, 1990). Feed samples were also analyzed for crude fibre content (method number 962.09). Digestible energy, and Ca and P were calculated from the ingredients of the experimental diets.

2.2. Nitrogen balance test with N-free ration

Twenty rabbits at the age of 16 weeks were used in a nitrogen balance trial with N-free ration. They were randomly divided into five groups (A–E). Before the test, all the rabbits were kept individually in cages and offered a diet with 160 g CP/kg and 10.4 MJ DE/kg for 7 days.

In the subsequent balance period, the rabbits were fed on a N-free diet, based on maize starch with the addition 140 g pure cellulose and 40 g premix (consisting of multi-vitamins and trace minerals) per kilogram. The daily allowance was 55, 45, 35, 25 and 0g/day for groups (Table 2).

The experimental period lasted for 6 days and excreta collection commenced on the 4th day. The collection methods were the same as described above.

2.3. Statistical analysis

Data from the N balance test and N-free diet trial were analyzed by two-ways and one-way ANOVA using MIXED procedure of SAS (1996), respectively. The relationships between different parameters were analyzed by linear regression (Snedecor and Cochran, 1989).

Table 2

The ex	perimental	design	for nitrogen	(N) balance with	N-free diet	NFD)
THC CA	permitticitui	acoign	ior merogen		bulunce with	it nee alee	1110,

Period	Groups						
	A	В	С	D	E		
Interim period							
Duration (day)	7	7	7	7	7		
Feed	Normal ^a	Normal	Normal	Normal	Normal		
Daily intake (g/day)	160	160	160	160	160		
Experimental period							
Duration (day)	6	6	6	6	6		
Feed	NFD	NFD	NFD	NFD	Fasting		
Daily intake (g/day)	55	45	35	25	0		
Days to start collection of excreta	4	4	4	4	After 40 h		
Duration of sampling (day)	3	3	3	3	40 h		

^a A diet with 160 g CP/kg and 10.4 MJ DE/kg.

3. Results and discussion

3.1. Nitrogen balance test

The results of N balance for rabbits over two periods are shown in Table 3. Net protein utilization (NPU) tended to increase with the increased levels of CP, but no significant difference was found among the five CP contents (P>0.05). Age had a significant influence on the NUP (P<0.001) that was lower during the growing period than during finishing period (P<0.01) possibly due to the fact that the mature rabbits can engage in coprophagy to use the non-protein nitrogenous compounds more efficiently than growing rabbits (Hoover and Heitmann, 1975). Because the contribution of microbial protein to young rabbits is small (Varela-Alvarez, 1991), the growing rabbits should be supplied with good-quality protein. On the other hand, the digestible capacity in the finishing rabbits should be higher than in the grower, resulting in a higher NPU in finishing period.

Table 3

Experimental period (age)	Designed CP (g/kg)	Initial weight (kg)	CP (g/kg)	Nitrogen balance (g/day)		NPU (g/kg)	
				Intake	Retention	NPU	Mean
Growing	120	2.1	115	2.8	1.2	440	445
period	140	2.1	137.5	3.0	1.2	405	
(6–12	160	2.1	159	3.8	1.7	454	
wks)	180	2.2	175	4.2	1.8	427	
	200	2.2	205	5.2	2.6	499	
Finishing	120	2.8	115	2.9	1.2	424	497
period	140	2.7	137.5	3.5	1.8	521	
(13-17	160	2.6	159	3.9	2.0	501	
wks)	180	2.7	175	4.3	2.2	501	
,	200	2.8	205	5.1	2.7	535	
S.E.M.		0.02		0.04	0.04	8.9	
Р	Age	<0.001		0.052	0.004	0.005	
	CP	0.193		< 0.001	< 0.001	0.061	
	$Age \times CP$	0.226		0.0097	0.255	0.210	

Content of crude protein (CP) in the diet, intake and retention of nitrogen (N), and net protein utilization (NPU) in WHBE rabbit at different ages.

3.2. The requirement of protein for maintenance estimated from N balance test

Taking the N retained in the body (NR) as the dependent variable and the N intake (NI) as the independent variable, the data were subject to linear regression. The results are shown in Fig. 1. The requirement of N for maintenance (Nm) may normally be estimated by extrapolating from the linear regression (McDonald et al., 2001) and is equivalent to NI where the relationship is extrapolated to zero N retention. The N requirement would increase with the increasing age and body weight, but no significant difference was found between the two periods when the Nm was express as the value per unit of metabolic body size (kg^{0.75}). The result was consistent with Miyasaki et al. (1997) who showed that the CP requirement expressed as gram per unit of metabolic body size remained constant with advancing age in broilers.

It was estimated that the requirement of Nm in WHBE rabbits was, on average, 485 mg/kg BW^{0.75}, which is equal to 3.03 g CP/kg BW^{0.75} per day. The value is lower than growing Angora wool-type rabbits (4.87 g CP/kg BW^{0.75}, Liu et al., 1991). The body weight of finishing rabbits was about 3.0 kg, so they need 6.91 g CP for maintenance. Given that the daily feed intake of a finishing rabbit was 150 g/day, the CP concentration of feed for maintenance is about 45.6 g/kg, which was lower than that recommended by NRC (1977) for meat-type rabbits (120 g/kg). The lower CP requirement for maintenance in WHBE rabbits than in wool-type and meat-type rabbits may be attributable to a lower basic metabolic rate in WHBE rabbits.

Assuming that the digestibility coefficient of dietary CP is 0.70, the equations in Fig. 1 may be converted to DCPI = 2.14 + 1.23CPR for growing rabbits, and DCPI = 2.11 + 1.03CPR for finishing rabbits, where DCPI and CPR are digestible CP intake and CP retention, as g/kg BW^{0.75} per day. From these two equations, the intercepts (2.14 and 2.11 g/kg BW^{0.75} per day) correspond to the DCP requirement for maintenance, and 0.813 (1/1.23) and 0.969 (1/1.03) are the amounts of dietary digestible CP used for growth that is retained in the bodies of growing and finishing rabbits, respectively. These constants are lower compared with those in the equation: DCPI = 2.88 + 1.78CPR for growing rabbit recommended by Fraga (1998), but resemble those of Spanish Giant rabbits (DCPI = 3.8 + 1.116CPR, de Blas et al., 1985).

3.3. Nitrogen balance test with N-free diet

The results for N balance test with N-free ration are shown in Table 4. When the rabbits are offered N-free diet, the total N excreted (TNE) may be considered as the requirement of Nm. It was found from



Fig. 1. Linear equation between nitrogen retained in the body (NR) and nitrogen intake (NI) during growing (♦) and finishing (▲) periods.

Groups	Body weight	Feed intake	N in excreta	N for maintena	nce
	kg	g/day	mg/day	mg/kgBW	mg/kgBW ^{0.75}
A	2.7	47.1	680	251	322
В	2.6	38.2	750	275	349
С	2.7	31.6	839	308	396
D	2.6	19.2	925	353	449
Е	2.6	0	1002	383	486
S.E.M.	0.05	0.46	26.2	13.4	15.5
Р	0.936	<0.001	0.010	0.046	0.029

The result for nitrogen (N) balance with the N-free ration and estimated N requirement for maintenance.

the trial that the lower the intake of N-free diet (FI), the greater the TNE. When the data were fitted to the linear regression, the following equations were obtained for the TNE (g/day) and FI (g/day): TNE = 1.022 - 0.007FI ($R^2 = 0.637$, P<0.01).

In fasting animals, some body protein might be catabolized to be used as a source of glucose via gluconeogenesis. In the animals offered N-free diet, the metabolic fecal N should be low due to low feed intake. Therefore, the TNE was mainly from the endogenous urinary N. The increased intake of N-free diet will help reduce the protein catabolism, resulting in less endogenous urinary N, and hence less TNE.

The rabbits in group E were fasted and the measured Nm was 486 mg/kg BW^{0.75}. This was in good agreement with the value obtained from the previous N balance test, confirming the result of N requirement. It implies that the extrapolation from the linear regression of NR on NI was valid, although none of the treatments were close to zero NR in the current study.

In summary, the requirement of Nm for WHBE rabbit is about 485 mg/kg BW^{0.75} per day. Further work is needed to study the requirement of other nutritional ingredients such as amino acids for this new strain of laboratory rabbit.

Acknowledgement

This work was supported in part by Zhejiang Provincial Department of Health, PR China (Grant no. 2006A094).

References

Amalia, E.Y., 2004. The laboratory rabbit: an animal model of atherosclerosis research. Lab Animals. 38, 246–256.

- AQSIQ (General Administration of Quality Supervision, Inception and Quarantine of the People's Republic of China), 2001a. Laboratory animals – standards monitoring for parasitology GB 14922.1-2001. Standards Press of China, Beijing.
- AQSIQ (General Administration of Quality Supervision, Inception and Quarantine of the People's Republic of China), 2001b. Laboratory animals – microbiological standards and monitoring GB14922.2-2001. Standards Press of China, Beijing.

Association of Official Analytical Chemist (AOAC), 1990. Official Methods of Analysis, 15th edition. AOAC, Arlington, VA, pp. 69–90.

Chen, M.L., Zhao, W.C., Ying, H.Z., Xu, X.P., Wang, D.J., 2005. RAPD analysis of the genetic specificity of WHBE rabbit. J. Zhejian Univ. (Agriculture & Life) 31 (4), 493–498 (in Chinese with English summary).

de Blas, J.C., María, J., Rodríguez, J.M., 1985. Units for feed evaluation and requirements for commercially grown rabbits. J. Anim. Sci. 60, 1021–1028.

Fraga, M.J., 1998. Protein requirements. In: de Blas, C., Wiseman, J. (Eds.), The Nutrition of the Rabbit. CABI Publishing, Wallingford, UK, pp. 113–143.

Hoover, W.H., Heitmann, R.N., 1975. Cercal nitrogen metabolism and amino acid absorption in the rabbit. J. Nutr. 105, 245–252.

Li, H., Wei, Y.X., 2002. Synthetical evaluation of dietary nutritional levels and commendatory feeding standard in rabbits. China Herbivores 2 (2), 38–40 (in Chinese).

Liu, S.M., Zhang, L., Wei, C.M., Chang, X., Lu, Y.X., Peng, D.H., 1991. The maintenance requirements of protein for Angora rabbits and the efficiency of digested protein by the rabbits. Acta Vet. Zootech. Sin. 22 (4), 323–326 (in Chinese with English summary).

Maerten, L., Luzi, F., De Groote, G., 1997. Effect of dietary protein and amino acids on the performance, carcass composition and N-excretion of growing rabbits. Ann. Zootech. 46, 255.

Maerten, L., 1992. Rabbit nutrition and feeding: a review of some developments. J. Appl. Rabbit Res. 15, 889.

McDonald, P., Edwards, R.A., Greenhalgh, J.F.D., 2001. Animal Nutrition, 6th edition. Longman/Wiley, New York.

Miyasaki, N., Takizawa, K., Hirata, T., Ueno, K., Kadowaki, M., Ishibashi, T., 1997. Effects of age on amino acid requirement for maintenance of broilers. Hokuriku J. Zootech. Sci. 74, 56–59.

Table 4

Varela-Alvarez, H., 1991. Feeding rabbits. In: Church, D.C. (Ed.), Livestock Feeds and Feeding, 3rd edition. Prentice Hall Inc., Englewood Cliffs, NJ, p. 454.

Wei, H., 1998. Medical Laboratory Animal Science. Science and Technology Press, Sichuang, PR China, p. 183.